

ZFP64 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19209b**Specification**

ZFP64 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9NTW7
Other Accession	NP_955459.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	72217
Antigen Region	525-551

ZFP64 Antibody (C-term) - Additional Information**Gene ID** 55734**Other Names**

Zinc finger protein 64 homolog, isoforms 3 and 4, Zfp-64, Zinc finger protein 338, ZFP64, ZNF338

Target/Specificity

This ZFP64 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 525-551 amino acids from the C-terminal region of human ZFP64.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZFP64 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZFP64 Antibody (C-term) - Protein Information**Name** ZFP64 ([HGNC:15940](#))**Function** May be involved in the regulation of mesenchymal cell differentiation through

transactivation of NOTCH1 target genes.

Cellular Location


Nucleus.

ZFP64 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZFP64 Antibody (C-term) - Images



Anti-ZFP64 Antibody (C-term) at 1:1000 dilution + human uterus lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 72 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ZFP64 Antibody (C-term) - Background

ZFP64 may be involved in transcriptional regulation.

ZFP64 Antibody (C-term) - References

Nicolas, E., et al. Eur. J. Hum. Genet. 18(10):1107-1113(2010) Davis, O.S., et al. Behav. Genet. (2010) In press : Aston, K.I., et al. J. Androl. 30(6):711-725(2009) Schymick, J.C., et al. Lancet Neurol 6(4):322-328(2007) Stelzl, U., et al. Cell 122(6):957-968(2005)