

Mouse Nod1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP19219c

Specification

Mouse Nod1 Antibody (Center) - Product Information

Application WB,E
Primary Accession Q8BHB0

Other Accession NP 001164478.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Mouse
Rabbit
Rabbit
Polyclonal
Rabbit IgG
107740
556-584

Mouse Nod1 Antibody (Center) - Additional Information

Gene ID 107607

Other Names

Nucleotide-binding oligomerization domain-containing protein 1, Caspase recruitment domain-containing protein 4, Nod1, Card4

Target/Specificity

This Mouse Nod1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 556-584 amino acids from the Central region of mouse Nod1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Nod1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Nod1 Antibody (Center) - Protein Information

Name Nod1 {ECO:0000303|PubMed:16211083, ECO:0000312|MGI:MGI:1341839}



Tel: 858.875.1900 Fax: 858.875.1999

Function Pattern recognition receptor (PRR) that detects bacterial peptidoglycan fragments and other danger signals and thus participates in both innate and adaptive immune responses (PubMed:12796777, PubMed:21715553). Specifically recognizes and binds gamma-D-glutamylmeso-diaminopimelic acid (iE-DAP), a dipeptide present in peptidoglycan of Gram-negative bacteria (PubMed:12796777, PubMed:16211083). Preferentially binds iE-DAP in tetrapeptide-containing muropeptides (MurNAc-TetraDAP or TetraDAP) (PubMed: 16211083). Ligand binding triggers oligomerization that facilitates the binding and subsequent activation of the proximal adapter receptor-interacting RIPK2 (By similarity). Following recruitment, RIPK2 undergoes 'Met-1'- (linear) and 'Lys-63'-linked polyubiquitination by E3 ubiquitin-protein ligases XIAP, BIRC2, BIRC3 and the LUBAC complex, becoming a scaffolding protein for downstream effectors, triggering activation of the NF- kappa-B and MAP kinases signaling (By similarity). This in turn leads to the transcriptional activation of hundreds of genes involved in immune response (By similarity). Also acts as a regulator of antiviral response elicited by dsRNA and the expression of RLR pathway members by targeting IFIH1 and TRAF3 to modulate the formation of IFIH1-MAVS and TRAF3-MAVS complexes leading to increased transcription of type I IFNs (By similarity). Also acts as a regulator of autophagy via its interaction with ATG16L1, possibly by recruiting ATG16L1 at the site of bacterial entry (PubMed: 19898471). Besides recognizing pathogens, also involved in the endoplasmic reticulum stress response: acts by sensing and binding to the cytosolic metabolite sphingosine-1-phosphate generated in response to endoplasmic reticulum stress, initiating an inflammation process that leads to activation of the NF-kappa-B and MAP kinases signaling (PubMed: 27007849). In addition, plays a role in insulin trafficking in beta cells in a cell-autonomous manner (PubMed:21715553, PubMed:31201384). Mechanistically, upon recognizing cognate ligands, NOD1 and RIPK2 localize to insulin vesicles where they recruit RAB1A to direct insulin trafficking through the cytoplasm (PubMed:31201384).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q9Y239}; Lipid-anchor {ECO:0000250|UniProtKB:Q9Y239}. Apical cell membrane {ECO:0000250|UniProtKB:Q9Y239}. Basolateral cell membrane {ECO:0000250|UniProtKB:Q9Y239}. Cytoplasm {ECO:0000250|UniProtKB:Q9Y239}. Note=Detected in the cytoplasm and at the cell membrane. Following bacterial infection, localizes to bacterial entry sites in the cell membrane. Recruited to the basolateral and apical membranes in polarized epithelial cells {ECO:0000250|UniProtKB:Q9Y239}

Tissue Location

Although ubiquitously expressed, NOD1 levels are more abundant in immune cells, the gastrointestinal tract, and adipose tissue.


Mouse Nod1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Nod1 Antibody (Center) - Images





Anti-Mouse Nod1 Antibody (Center) at 1:1000 dilution + mouse spleen lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 108 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse Nod1 Antibody (Center) - Background

Enhances caspase-9-mediated apoptosis. Induces NF-kappa-B activity via RIPK2 and IKK-gamma. Confers responsiveness to intracellular bacterial lipopolysaccharides (LPS). Forms an intracellular sensing system along with ARHGEF2 for the detection of microbial effectors during cell invasion by pathogens (By similarity).

Mouse Nod1 Antibody (Center) - References

Moreno, L., et al. Br. J. Pharmacol. 160(8):1997-2007(2010) lyer, J.K., et al. Infect. Immun. 78(6):2418-2428(2010) Watanabe, T., et al. J. Clin. Invest. 120(5):1645-1662(2010) Toma, C., et al. J. Immunol. 184(9):5287-5297(2010) Shigeoka, A.A., et al. J. Immunol. 184(5):2297-2304(2010)