

KCNE2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19267b**Specification**

KCNE2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9Y6J6
Other Accession	NP_751951.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	14472
Antigen Region	79-107

KCNE2 Antibody (C-term) - Additional Information**Gene ID** 9992**Other Names**

Potassium voltage-gated channel subfamily E member 2, MinK-related peptide 1, Minimum potassium ion channel-related peptide 1, Potassium channel subunit beta MiRP1, KCNE2

Target/Specificity

This KCNE2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 79-107 amino acids from the C-terminal region of human KCNE2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KCNE2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

KCNE2 Antibody (C-term) - Protein Information**Name** KCNE2**Function** Ancillary protein that assembles as a beta subunit with a voltage-gated potassium

channel complex of pore-forming alpha subunits. Modulates the gating kinetics and enhances stability of the channel complex. Assembled with KCNB1 modulates the gating characteristics of the delayed rectifier voltage-dependent potassium channel KCNB1. Associated with KCNH2/HERG is proposed to form the rapidly activating component of the delayed rectifying potassium current in heart (IKr). May associate with KCNQ2 and/or KCNQ3 and modulate the native M-type current. May associate with HCN1 and HCN2 and increase potassium current. Interacts with KCNQ1; forms a heterooligomer complex leading to currents with an apparently instantaneous activation, a rapid deactivation process and a linear current-voltage relationship and decreases the amplitude of the outward current (PubMed:[11101505](#)). KCNQ1-KCNE2 channel associates with Na(+)-coupled myo-inositol symporter in the apical membrane of choroid plexus epithelium and regulates the myo-inositol gradient between blood and cerebrospinal fluid with an impact on neuron excitability.

Cellular Location

Cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P63161} Apical cell membrane {ECO:0000250|UniProtKB:Q9D808}; Single-pass membrane protein.
Note=Colocalizes with KCNB1 at the plasma membrane. {ECO:0000250|UniProtKB:P63161}

Tissue Location


Highly expressed in brain, heart, skeletal muscle, pancreas, placenta, kidney, colon and thymus. A small but significant expression is found in liver, ovary, testis, prostate, small intestine and leukocytes. Very low expression, nearly undetectable, in lung and spleen.

KCNE2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KCNE2 Antibody (C-term) - Images



KCNE2 Antibody (C-term)(Cat. #AP19267b) western blot analysis in A549 cell line lysates

(35ug/lane). This demonstrates the KCNE2 antibody detected the KCNE2 protein (arrow).

KCNE2 Antibody (C-term) - Background

Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, *isk*-related subfamily. This member is a small integral membrane subunit that assembles with the KCNH2 gene product, a pore-forming protein, to alter its function. This gene is expressed in heart and muscle and the gene mutations are associated with cardiac arrhythmia.

KCNE2 Antibody (C-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Albert, C.M., et al. Circ Arrhythm Electrophysiol 3(3):222-229(2010)
Subbiah, R.N., et al. Can J Cardiol 26(4):208-212(2010)
Tam, G.W., et al. Biochem. Soc. Trans. 38(2):445-451(2010)
Roepke, T.K., et al. PLoS ONE 5 (7), E11451 (2010) :