

APC Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19274c**Specification**

APC Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P25054
Other Accession	NP_000029.2
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1148-1176

APC Antibody (Center) - Additional Information**Gene ID** 324**Other Names**

Adenomatous polyposis coli protein, Protein APC, Deleted in polyposis 25, APC, DP25

Target/Specificity

This APC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1148-1176 amino acids from the Central region of human APC.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

APC Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

APC Antibody (Center) - Protein Information**Name** APC ([HGNC:583](#))**Synonyms** DP2.5

Function Tumor suppressor. Promotes rapid degradation of CTNNB1 and participates in Wnt signaling as a negative regulator. APC activity is correlated with its phosphorylation state. Activates the GEF activity of SPATA13 and ARHGEF4. Plays a role in hepatocyte growth factor (HGF)- induced cell migration. Required for MMP9 up-regulation via the JNK signaling pathway in colorectal tumor cells. Associates with both microtubules and actin filaments, components of the cytoskeleton (PubMed:[17293347](#)). Plays a role in mediating the organization of F- actin into ordered bundles (PubMed:[17293347](#)). Functions downstream of Rho GTPases and DIAPH1 to selectively stabilize microtubules (By similarity). Acts as a mediator of ERBB2-dependent stabilization of microtubules at the cell cortex. It is required for the localization of MACF1 to the cell membrane and this localization of MACF1 is critical for its function in microtubule stabilization.

Cellular Location

Cell junction, adherens junction. Cytoplasm, cytoskeleton. Cell projection, lamellipodium. Cell projection, ruffle membrane. Cytoplasm. Cell membrane. Note=Associated with the microtubule network at the growing distal tip of microtubules (PubMed:19632184) MAPRE1 may be required for targeting to the growing microtubule plus ends (PubMed:19632184). Accumulates in the lamellipodium and ruffle membrane in response to hepatocyte growth factor (HGF) treatment (PubMed:19151759). The MEMO1-RHOA-DIAPH1 signaling pathway controls localization of the phosphorylated form to the cell membrane (PubMed:20937854).

Tissue Location


Expressed in a variety of tissues: brain, small intestine, colon, thymus, skeletal muscle, heart, prostate, lung, spleen, ovary, testis kidney, placenta, blood and liver (PubMed:21643010, PubMed:27217144). Isoform 1A: Very strongly expressed in brain but has relatively low expression levels in other tissues (PubMed:19527921, PubMed:21643010, PubMed:27217144). Isoform 1B: Predominant form in all tissues except for brain, including gastric mucosa and blood (PubMed:19527921, PubMed:21643010, PubMed:27217144)


APC Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

APC Antibody (Center) - Images





APC Antibody (Center)(Cat. #AP19274c) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the APC antibody detected the APC protein (arrow).

APC Antibody (Center) - Background

This gene encodes a tumor suppressor protein that acts as an antagonist of the Wnt signaling pathway. It is also involved in other processes including cell migration and adhesion, transcriptional activation, and apoptosis. Defects in this gene cause familial adenomatous polyposis (FAP), an autosomal dominant pre-malignant disease that usually progresses to malignancy. Disease-associated mutations tend to be clustered in a small region designated the mutation cluster region (MCR) and result in a truncated protein product.

APC Antibody (Center) - References

Chung, K.H., et al. J Neurosurg Pediatr 6(4):372-376(2010)
Rai, K., et al. Cell 142(6):930-942(2010)
Poulogiannis, G., et al. Proc. Natl. Acad. Sci. U.S.A. 107(34):15145-15150(2010)
Jaulin, F., et al. J. Cell Biol. 190(3):443-460(2010)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)