

Mouse Rcvrn Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19308a**Specification**

Mouse Rcvrn Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P34057
Other Accession	NP_033064.1
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	23407
Antigen Region	39-67

Mouse Rcvrn Antibody (N-term) - Additional Information**Gene ID** 19674**Other Names**

Recoverin, 23 kDa photoreceptor cell-specific protein, Cancer-associated retinopathy protein, Protein CAR, Rcvrn, Rcv1

Target/Specificity

This Mouse Rcvrn antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 39-67 amino acids from the N-terminal region of mouse Rcvrn.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Rcvrn Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Rcvrn Antibody (N-term) - Protein Information**Name** Rcvrn

Synonyms Rcv1

Function Acts as a calcium sensor and regulates phototransduction of cone and rod photoreceptor cells (By similarity). Modulates light sensitivity of cone photoreceptor in dark and dim conditions (PubMed:[25673692](#)). In response to high Ca^{2+} levels induced by low light levels, prolongs RHO/rhodopsin activation in rod photoreceptor cells by binding to and inhibiting GRK1-mediated phosphorylation of RHO/rhodopsin (By similarity). Plays a role in scotopic vision/enhances vision in dim light by enhancing signal transfer between rod photoreceptors and rod bipolar cells (PubMed:[15882641](#)). Improves rod photoreceptor sensitivity in dim light and mediates response of rod photoreceptors to facilitate detection of change and motion in bright light (PubMed:[29435986](#)).

Cellular Location

Photoreceptor inner segment. Cell projection, cilium, photoreceptor outer segment Photoreceptor outer segment membrane {ECO:0000250|UniProtKB:P21457}; Lipid-anchor {ECO:0000250|UniProtKB:P21457}; Cytoplasmic side {ECO:0000250|UniProtKB:P21457}. Perikaryon. Note=Primarily expressed in the inner segments of light-adapted rod photoreceptors, approximately 10% of which translocates from photoreceptor outer segments upon light stimulation (PubMed:15961391). Targeting of myristoylated protein to rod photoreceptor outer segments is calcium dependent (By similarity) {ECO:0000250|UniProtKB:P21457, ECO:0000269|PubMed:15961391}

Tissue Location


Expressed in rod photoreceptors in the retina (at protein level).

Mouse Rcvrn Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Rcvrn Antibody (N-term) - Images



Mouse Rcvrn Antibody (N-term)(Cat. #AP19308a) western blot analysis in mouse kidney tissue lysates (35ug/lane). This demonstrates the Rcvrn antibody detected the Rcvrn protein (arrow).

Mouse Rcvrn Antibody (N-term) - Background

Rcvrn seems to be implicated in the pathway from retinal rod guanylate cyclase to rhodopsin. May be involved in the inhibition of the phosphorylation of rhodopsin in a calcium-dependent manner. The calcium-bound recoverin prolongs the photoresponse.

Mouse Rcvrn Antibody (N-term) - References

Jin, K., et al. J. Neurosci. 30(36):11902-11916(2010)
Georgi, S.A., et al. J. Neurosci. 30(11):4048-4061(2010)
Sanuki, R., et al. FEBS Lett. 584(4):753-758(2010)
Brzezinski, J.A. IV, et al. Development 137(4):619-629(2010)
Chen, C.K., et al. J. Neurosci. 30(4):1213-1220(2010)