

### DDX41 Antibody(N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP19450a

### **Specification**

## DDX41 Antibody(N-term) - Product Information

Application WB,E
Primary Accession Q9UJV9

Other Accession <u>Q91VN6</u>, <u>NP 057306.2</u>

Reactivity
Predicted
Host
Clonality
Isotype
Calculated MW
Antigen Region

Human
Mouse
Rabbit
Polyclonal
Rabbit IgG
141-169

## DDX41 Antibody(N-term) - Additional Information

#### **Gene ID 51428**

### **Other Names**

Probable ATP-dependent RNA helicase DDX41, DEAD box protein 41, DEAD box protein abstrakt homolog, DDX41, ABS

### Target/Specificity

This DDX41 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 141-169 amino acids from the N-terminal region of human DDX41.

#### **Dilution**

WB~~1:1000

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

DDX41 Antibody(N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# DDX41 Antibody(N-term) - Protein Information

#### Name DDX41





**Synonyms** ABS

**Function** Probable ATP-dependent RNA helicase. Is required during post- transcriptional gene expression. May be involved in pre-mRNA splicing.


**Cellular Location** Nucleus.

### DDX41 Antibody(N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


## DDX41 Antibody(N-term) - Images



DDX41 Antibody (N-term) (Cat. #AP19450a) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the DDX41 antibody detected the DDX41 protein (arrow).

# DDX41 Antibody(N-term) - Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of the DEAD box protein family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a member of this family. The function of this member has not been determined. Based on studies in Drosophila, the abstrakt gene is widely required during post-transcriptional gene expression.





# DDX41 Antibody(N-term) - References

Wu, C., et al. Proteomics 7(11):1775-1785(2007)

Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007):

Olsen, J.V., et al. Cell 127(3):635-648(2006)

Olsen, J.V., et al. Cell 127(3):635-648(2006)

Nousiainen, M., et al. Proc. Natl. Acad. Sci. U.S.A. 103(14):5391-5396(2006)