

**Mouse Dab2 Antibody(C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP19696b****Specification**

---

**Mouse Dab2 Antibody(C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P98078</a>
Other Accession	<a href="#">O88797</a> , <a href="#">NP_001008702.1</a> , <a href="#">NP_001032994.1</a> , <a href="#">NP_075607.2</a>
Reactivity	Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	82312
Antigen Region	624-650

**Mouse Dab2 Antibody(C-term) - Additional Information****Gene ID** 13132**Other Names**

Disabled homolog 2, DOC-2, Mitogen-responsive phosphoprotein, Dab2, Doc2

**Target/Specificity**

This Mouse Dab2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 624-650 amino acids from the C-terminal region of mouse Dab2.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Mouse Dab2 Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**Mouse Dab2 Antibody(C-term) - Protein Information****Name** Dab2

## Synonyms Doc2

**Function** Adapter protein that functions as a clathrin-associated sorting protein (CLASP) required for clathrin-mediated endocytosis of selected cargo proteins. Can bind and assemble clathrin, and binds simultaneously to phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P<sub>2</sub>) and cargos containing non-phosphorylated NPXY internalization motifs, such as the LDL receptor, to recruit them to clathrin-coated pits. Can function in clathrin-mediated endocytosis independently of the AP-2 complex. Involved in endocytosis of integrin beta-1; this function seems to be redundant with the AP-2 complex and seems to require DAB2 binding to endocytosis accessory EH domain-containing proteins such as EPS15, EPS15L1 and ITSN1. Involved in endocytosis of cystic fibrosis transmembrane conductance regulator/CFTR. Isoform p96 is involved in endocytosis of megalin/LRP2 lipoprotein receptor during embryonal development. Required for recycling of the TGF-beta receptor. Isoform p67 is not involved in LDL receptor endocytosis. Involved in CFTR trafficking to the late endosome. Involved in several receptor-mediated signaling pathways. Involved in TGF-beta receptor signaling and facilitates phosphorylation of the signal transducer SMAD2. Mediates TGF-beta-stimulated JNK activation. May inhibit the canonical Wnt/beta-catenin signaling pathway by stabilizing the beta-catenin destruction complex through a competing association with axin preventing its dephosphorylation through protein phosphatase 1 (PP1). Sequesters LRP6 towards clathrin-mediated endocytosis, leading to inhibition of Wnt/beta-catenin signaling. May activate non-canonical Wnt signaling. In cell surface growth factor/Ras signaling pathways proposed to inhibit ERK activation by interrupting the binding of GRB2 to SOS1 and to inhibit SRC by preventing its activating phosphorylation at 'Tyr-419'. Proposed to be involved in modulation of androgen receptor (AR) signaling mediated by SRC activation; seems to compete with AR for interaction with SRC. Plays a role in the CSF-1 signal transduction pathway. Plays a role in cellular differentiation. Involved in cell positioning and formation of visceral endoderm (VE) during embryogenesis and proposed to be required in the VE to respond to Nodal signaling coming from the epiblast. Required for the epithelial to mesenchymal transition, a process necessary for proper embryonic development. May be involved in myeloid cell differentiation and can induce macrophage adhesion and spreading. Isoform p67 may be involved in transcriptional regulation. May act as a tumor suppressor.

## Cellular Location

Cytoplasmic vesicle, clathrin-coated vesicle membrane. Membrane, clathrin-coated pit.  
Note=Colocalizes with large insert-containing isoforms of MYO6 at clathrin-coated pits/vesicles During mitosis is progressively displaced from the membrane and translocated to the cytoplasm (By similarity). [Isoform p67]: Cytoplasm. Nucleus Note=Diffuse localization in the cytoplasm; does not localize to clathrin-coated pits

## Tissue Location


Isoform p96 and isoform p67 are expressed in adult kidney and fibroblasts with isoform p96 being the predominant form Isoform p67 is the predominant isoform expressed in embryonic visceral endoderm.

## Mouse Dab2 Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Mouse Dab2 Antibody(C-term) - Images



Mouse Dab2 Antibody (C-term) (Cat. #AP19696b) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the Mouse Dab2 antibody detected the Mouse Dab2 protein (arrow).

### Mouse Dab2 Antibody(C-term) - Background

Component of the CSF-1 signal transduction pathway.

### Mouse Dab2 Antibody(C-term) - References

- Collaco, A., et al. J. Biol. Chem. 285(22):17177-17187(2010)
- Chaudhury, A., et al. Nat. Cell Biol. 12(3):286-293(2010)
- Hasumi, Y., et al. Proc. Natl. Acad. Sci. U.S.A. 106(44):18722-18727(2009)
- Jain, N., et al. J. Immunol. 183(7):4192-4196(2009)
- Jiang, Y., et al. Oncogene 28(33):2999-3007(2009)