

PTBP1 Antibody(N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19710a

Specification

PTBP1 Antibody(N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P26599
Other Accession	Q8WN55 , NP_002810.1 , NP_114367.1 , NP_114368.1 , NP_787041.1
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	26-55

PTBP1 Antibody(N-term) - Additional Information

Gene ID 5725

Other Names

Polypyrimidine tract-binding protein 1, PTB, 57 kDa RNA-binding protein PPTB-1, Heterogeneous nuclear ribonucleoprotein I, hnRNP I, PTBP1, PTB

Target/Specificity

This PTBP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 26-55 amino acids from the N-terminal region of human PTBP1.

Dilution

WB~~1:1000
IHC-P~~1:4000
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PTBP1 Antibody(N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PTBP1 Antibody(N-term) - Protein Information

Name PTBP1

Synonyms PTB

Function Plays a role in pre-mRNA splicing and in the regulation of alternative splicing events. Activates exon skipping of its own pre- mRNA during muscle cell differentiation. Binds to the polypyrimidine tract of introns. May promote RNA looping when bound to two separate polypyrimidine tracts in the same pre-mRNA. May promote the binding of U2 snRNP to pre-mRNA. Cooperates with RAVER1 to modulate switching between mutually exclusive exons during maturation of the TPM1 pre- mRNA. Represses the splicing of MAPT/Tau exon 10 (PubMed:[15009664](#)). Binds to polypyrimidine-rich controlling element (PCE) of CFTR and promotes exon skipping of CFTR exon 9, thereby antagonizing TIA1 and its role in exon inclusion of CFTR exon 9 (PubMed:[14966131](#)). Plays a role in the splicing of pyruvate kinase PKM by binding repressively to a polypyrimidine tract flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:[20010808](#)). In case of infection by picornaviruses, binds to the viral internal ribosome entry site (IRES) and stimulates the IRES- mediated translation (PubMed:[21518806](#)).

Cellular Location


Nucleus.

PTBP1 Antibody(N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PTBP1 Antibody(N-term) - Images



PTBP1 Antibody (N-term) (Cat. #AP19710a) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the PTBP1 antibody detected the PTBP1 protein (arrow).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue using AP19710a. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems was used as the secondary antibody.

PTBP1 Antibody(N-term) - Background

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA-binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has four repeats of quasi-RNA recognition motif (RRM) domains that bind RNAs. This protein binds to the intronic polypyrimidine tracts that requires pre-mRNA splicing and acts via the protein degradation ubiquitin-proteasome pathway. It may also promote the binding of U2 snRNP to pre-mRNAs. This protein is localized in the nucleoplasm and it is also detected in the perinucleolar structure. Alternatively spliced transcript variants encoding different isoforms have been described.

PTBP1 Antibody(N-term) - References

Kanda, T., et al. J. Viral Hepat. 17(9):618-623(2010)
Cobbold, L.C., et al. Oncogene 29(19):2884-2891(2010)
Verma, B., et al. J. Gen. Virol. 91 (PT 5), 1245-1255 (2010) :
Maynard, C.M., et al. J. Mol. Biol. 397(1):260-277(2010)
Xue, Y., et al. Mol. Cell 36(6):996-1006(2009)