

HDBP2(HDRF-2) Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP1979c

Specification

HDBP2(HDRF-2) Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	<u>Q9H8N7</u>
Other Accession	<u>NP_061130</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
•	

HDBP2(HDRF-2) Antibody (Center) - Additional Information

Gene ID 55893

Other Names

Zinc finger protein 395, HD-regulating factor 2, HDRF-2, Huntington disease gene regulatory region-binding protein 2, HD gene regulatory region-binding protein 2, HDBP-2, Papillomavirus regulatory factor 1, PRF-1, Papillomavirus-binding factor, ZNF395, HDBP2, PBF

Target/Specificity

This HDBP2(HDRF-2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 141-170 amino acids from the Central region of human HDBP2(HDRF-2).

Dilution WB~~1:1000 IHC-P~~1:50~100 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HDBP2(HDRF-2) Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

HDBP2(HDRF-2) Antibody (Center) - Protein Information



Name ZNF395

Synonyms HDBP2, PBF

Function Plays a role in papillomavirus genes transcription.

Cellular Location Cytoplasm. Nucleus. Note=May shuttle between nucleus and cytoplasm


Tissue Location Widely expressed.

HDBP2(HDRF-2) Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

HDBP2(HDRF-2) Antibody (Center) - Images



Western blot analysis of anti-HDBP2(HDRF-2) Antibody (Center) (Cat. #AP1979c) in Y79 cell line lysates (35ug/lane). HDBP2(arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human Hepatocellular carcinoma reacted with anti-HDBP2(HDRF-2) Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

HDBP2(HDRF-2) Antibody (Center) - Background

Papillomavirus Regulatory Factor 1/HDBP2 is a novel transcription factor shuttling between nucleus and cytoplasm and binds to the specific GCCGGCG, which is an essential cis-element for Huntington's disease gene expression.

HDBP2(HDRF-2) Antibody (Center) - References

Tanaka, K., J. Biol. Chem. 279 (8), 7275-7286 (2004)