

TAF11 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19984a

Specification

TAF11 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q15544
Other Accession	NP_005634.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	23307
Antigen Region	1-30

TAF11 Antibody (N-term) - Additional Information

Gene ID 6882

Other Names

Transcription initiation factor TFIID subunit 11, TFIID subunit p30-beta, Transcription initiation factor TFIID 28 kDa subunit, TAF(II)28, TAFII-28, TAFII28, TAF11, TAF2I

Target/Specificity

This TAF11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human TAF11.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TAF11 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TAF11 Antibody (N-term) - Protein Information

Name TAF11

Synonyms TAF2I

Function The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed:[33795473](#)). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed:[33795473](#)). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed:[33795473](#)). TAF11, together with TAF13 and TBP, play key roles during promoter binding by the TFIID and TFIIA transcription factor complexes (PubMed:[33795473](#)).

Cellular Location


Nucleus.

TAF11 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TAF11 Antibody (N-term) - Images



TAF11 Antibody (N-term) (Cat. #AP19984a) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the TAF11 antibody detected the TAF11 protein (arrow).

TAF11 Antibody (N-term) - Background

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory

signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a small subunit of TFIID that is present in all TFIID complexes and interacts with TBP. This subunit also interacts with another small subunit, TAF13, to form a heterodimer with a structure similar to the histone core structure.

TAF11 Antibody (N-term) - References

Matsuoka, S., et al. Science 316(5828):1160-1166(2007)
Mungall, A.J., et al. Nature 425(6960):805-811(2003)
Guermah, M., et al. Mol. Cell 12(4):991-1001(2003)
Mengus, G., et al. J. Biol. Chem. 275(14):10064-10071(2000)
Birck, C., et al. Cell 94(2):239-249(1998)