

BMPR1A Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2004a

Specification

BMPR1A Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P36894
Other Accession	Q78EA7 , P36895 , NP_004320
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	30-60

BMPR1A Antibody (N-term) - Additional Information

Gene ID 657

Other Names

Bone morphogenetic protein receptor type-1A, BMP type-1A receptor, BMPR-1A, Activin receptor-like kinase 3, ALK-3, Serine/threonine-protein kinase receptor R5, SKR5, CD292, BMPR1A, ACVRLK3, ALK3

Target/Specificity

This BMPR1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 30-60 amino acids from the N-terminal region of human BMPR1A.

Dilution

WB~~1:1000
IHC-P~~1:50~100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BMPR1A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of lysate from rat uterus tissue lysate, using BMPR1A Antibody (E45)(Cat. #AP2004a). AP2004a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.



Western blot analysis of anti-BMPR1A Pab (Cat. #ap2004a) in Saos-2 cell lysate. BMPR1A (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

BMPR1A Antibody (N-term) - Protein Information

Name BMPR1A

Synonyms ACVRLK3, ALK3

Function

On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. Receptor for BMP2, BMP4, GDF5 and GDF6. Positively regulates chondrocyte differentiation through GDF5 interaction. Mediates induction of adipogenesis by GDF6.

Cellular Location

Cell membrane
{ECO:0000250|UniProtKB:P36898};
Single-pass type I membrane protein. Cell surface {ECO:0000250|UniProtKB:P36895}


Tissue Location

Highly expressed in skeletal muscle.


BMPR1A Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Western blot analysis of lysate from mouse kidney tissue, using BMPR1A Antibody (E45)(Cat. #AP2004a). AP2004a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

BMPR1A Antibody (N-term) - Background

The bone morphogenetic protein (BMP) receptors are a family of transmembrane serine/threonine kinases that include the type I receptors BMPR1A and BMPR1B and the type II receptor BMPR2. These receptors are also closely related to the activin receptors, ACVR1 and ACVR2. The ligands of these receptors are members of the TGF-beta superfamily. TGF-betas and activins transduce their signals through the formation of heteromeric complexes with 2 different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II

receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding.

BMPR1A Antibody (N-term) - References

Kan, L. et al. Stem Cells. January; 27(1): 150-156(2009). Waite, K.A., et al., Hum. Mol. Genet. 12(6):679-684 (2003). Zhou, X.P., et al., Am. J. Hum. Genet. 69(4):704-711 (2001). Astrom, A.K., et al., Mamm. Genome 10(3):299-302 (1999). ten Dijke, P., et al., Oncogene 8(10):2879-2887 (1993). Ide, H., et al., Cytogenet. Cell Genet. 81 (3-4), 285-286 (1998).

BMPR1A Antibody (N-term) - Citations

- [Targeted Knockdown of Bone Morphogenetic Protein Signaling within Neural Progenitors Protects the Brain and Improves Motor Function following Postnatal Hypoxia-Ischemia.](#)
- [BMP receptor 1A regulates development of hypothalamic circuits critical for feeding behavior.](#)
- [Differential effects of BMP signaling on parvalbumin and somatostatin interneuron differentiation.](#)
- [Dysregulation of local stem/progenitor cells as a common cellular mechanism for heterotopic ossification.](#)
- [BMP inhibition enhances axonal growth and functional recovery after spinal cord injury.](#)