

Mouse Sike1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20223a

Specification

Mouse Sike1 Antibody (N-term) - Product Information

Application WB,E
Primary Accession Q9CPR7

Other Accession Q5FWT9, NP_079955.1

Reactivity
Predicted
Rat
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Mouse Sike1 Antibody (N-term) - Additional Information

Gene ID 66641

Other Names

Suppressor of IKBKE 1, Suppressor of IKK-epsilon, Sike1

Target/Specificity

This Mouse Sike1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 36-63 amino acids from the N-terminal region of mouse Sike1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Sike1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Sike1 Antibody (N-term) - Protein Information

Name Sike1

Function Physiological suppressor of IKK-epsilon and TBK1 that plays an inhibitory role in virus-



and TLR3-triggered IRF3. Inhibits TLR3- mediated activation of interferon-stimulated response elements (ISRE) and the IFN-beta promoter. May act by disrupting the interactions of IKBKE or TBK1 with TICAM1/TRIF, IRF3 and RIGI. Does not inhibit NF- kappa-B activation pathways (By similarity).


Cellular Location Cytoplasm.

Mouse Sike1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Mouse Sike1 Antibody (N-term) - Images



Anti-MOUSE Sike1 Antibody (N-term) at 1:1000 dilution + mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse Sike1 Antibody (N-term) - Background

Physiological suppressor of IKK-epsilon and TBK1 that plays an inhibitory role in virus-and TLR3-triggered IRF3. Inhibits TLR3-mediated activation of interferon-stimulated response elements (ISRE) and the IFN-beta promoter. May act by disrupting the interactions of IKBKE or TBK1 with TICAM1/TRIF, IRF3 and DDX58/RIG-I. Does not inhibit NF-kappa-B activation pathways (By similarity).

Mouse Sike1 Antibody (N-term) - References





Katayama, S., et al. Science 309(5740):1564-1566(2005) Carninci, P., et al. Science 309(5740):1559-1563(2005)