10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SNX12 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20307c

Specification

SNX12 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	Q9UMY4
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**18885**
Antigen Region	**84-112**

SNX12 Antibody (Center) - Additional Information

Gene ID 29934

Other Names
Sorting nexin-12, SNX12

Target/Specificity
This SNX12 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 84-112 amino acids from the Central region of human SNX12.

Dilution
WB~~1:1000
E~~Use at an assay dependent concentration.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
SNX12 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SNX12 Antibody (Center) - Protein Information

Name SNX12

Function May be involved in several stages of intracellular trafficking.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Cellular Location
Membrane; Peripheral membrane protein; Cytoplasmic side

SNX12 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SNX12 Antibody (Center) - Images



SNX12 Antibody (Center) (Cat. #AP20307c) western blot analysis in A2058 cell line lysates (35ug/lane).This demonstrates the SNX12 antibody detected the SNX12 protein (arrow).

SNX12 Antibody (Center) - Background

May be involved in several stages of intracellular trafficking (By similarity).