

**RBPMs Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20354b****Specification**

---

**RBPMs Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O93062</a>
Other Accession	<a href="#">O9WVB0</a> , <a href="#">A0A8I6G705</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21802
Antigen Region	166-193

**RBPMs Antibody (C-term) - Additional Information****Gene ID** 11030**Other Names**

RNA-binding protein with multiple splicing, RBP-MS, Heart and RRM expressed sequence, Hermes, RBPMs, HERMES

**Target/Specificity**

This RBPMs antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 166-193 amino acids from the C-terminal region of human RBPMs.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RBPMs Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**RBPMs Antibody (C-term) - Protein Information****Name** RBPMs ([HGNC:19097](#))

## Synonyms HERMES

**Function** [Isoform A]: RNA binding protein that mediates the regulation of pre-mRNA alternative splicing (AS) (PubMed:[24860013](#), PubMed:[26347403](#)). Acts either as activator (FLNB, HSPG2, LIPA1, MYOCD, PTPRF and PPFIBP1) or repressor (TPM1, ACTN1, ITGA7, PIEZO1, LSM14B, MBNL1 and MBML2) of splicing events on specific pre-mRNA targets (By similarity). Together with RNA binding proteins RBFOX2 and MBNL1/2, activates a splicing program associated with differentiated contractile vascular smooth muscle cells (SMC) by regulating AS of numerous pre- mRNA involved in actin cytoskeleton and focal adhesion machineries, suggesting a role in promoting a cell differentiated state (By similarity). Binds to introns, exons and 3'-UTR associated with tandem CAC trinucleotide motifs separated by a variable spacer region, at a minimum as a dimer. The minimal length of RNA required for RBPMS- binding tandem CAC motifs is 15 nt, with spacing ranging from 1 to 9 nt. Can also bind to CA dinucleotide repeats (PubMed:[24860013](#), PubMed:[26347403](#)). Mediates repression of TPM1 exon 3 by binding to CAC tandem repeats in the flanking intronic regions, followed by higher- order oligomerization and heterotypic interactions with other splicing regulators including MBNL1 and RBFOX2, which prevents assembly of ATP- dependent splicing complexes (By similarity).

## Cellular Location

Nucleus. Cytoplasm. Cytoplasm, Stress granule. Cytoplasm, P-body. Note=Localized to cytoplasmic stress granules after oxidative stress (PubMed:24860013). Translocates into cytoplasmic stress granules that probably corresponds to P-bodies in response to oxidative stress (PubMed:26347403)

## Tissue Location


Ubiquitously expressed, at various levels depending on the isoform and the tissue (PubMed:8855282). Strongly expressed in the heart, prostate, small intestine, large intestine, and ovary; moderately expressed in the placenta, lung, liver, kidney, pancreas, and testis; and poorly expressed in the skeletal muscle, spleen, thymus and peripheral leukocytes (PubMed:8855282)


## RBPMS Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## RBPMS Antibody (C-term) - Images





RBPMs Antibody (C-term) (Cat. #AP20354b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the RBPMs antibody detected the RBPMs protein (arrow).

#### **RBPMs Antibody (C-term) - Background**

Acts as a coactivator of transcriptional activity. Required to increase TGF $\beta$ 1/Smad-mediated transactivation. Acts through SMAD2, SMAD3 and SMAD4 to increase transcriptional activity. Increases phosphorylation of SMAD2 and SMAD3 on their C-terminal SSXS motif, possibly through recruitment of TGF $\beta$ 1. Promotes the nuclear accumulation of SMAD2, SMAD3 and SMAD4 proteins. Binds to poly(A) RNA.