

TMX1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20365b

Specification

TMX1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q9H3N1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	31791
Antigen Region	239-267

TMX1 Antibody (C-term) - Additional Information

Gene ID 81542

Other Names

Thioredoxin-related transmembrane protein 1, Thioredoxin domain-containing protein 1, Transmembrane Trx-related protein, TMX1, TMX, TXNDC, TXNDC1

Target/Specificity

This TMX1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 239-267 amino acids from the C-terminal region of human TMX1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TMX1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TMX1 Antibody (C-term) - Protein Information

Name TMX1 {ECO:0000303|PubMed:37648867, ECO:0000312|HGNC:HGNC:15487}

Function Thioredoxin domain-containing protein that participates in various redox reactions

through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions (PubMed:[11152479](#), PubMed:[37648867](#)). Acts as a key inhibitor of the alternative triglyceride biosynthesis pathway by inhibiting the activity of TMEM68/DIESL at the endoplasmic reticulum, thereby restricting accumulation of triacylglycerol (PubMed:[37648867](#)). The alternative triglyceride biosynthesis pathway mediates formation of triacylglycerol from diacylglycerol and membrane phospholipids (PubMed:[37648867](#)). Acts as a protein disulfide isomerase by catalyzing formation or reduction of disulfide bonds (PubMed:[22228764](#), PubMed:[29932915](#)). Specifically mediates formation of disulfide bonds of transmembrane proteins at the endoplasmic reticulum membrane (PubMed:[22228764](#)). Involved in endoplasmic reticulum-associated degradation (ERAD) via its protein disulfide isomerase activity by acting on folding-defective polypeptides at the endoplasmic reticulum membrane (PubMed:[29932915](#)). Acts as a negative regulator of platelet aggregation following secretion in the extracellular space (PubMed:[30425049](#)). Acts as a regulator of endoplasmic reticulum-mitochondria contact sites via its ability to regulate redox signals (PubMed:[27502484](#), PubMed:[31304984](#)). Regulates endoplasmic reticulum- mitochondria Ca(2+) flux (PubMed:[27502484](#)).

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Mitochondrion membrane; Single-pass type I membrane protein. Secreted. Note=Predominantly found in the endoplasmic reticulum (PubMed:11152479). Secreted in the extracellular space following thrombin stimulation (PubMed:30425049). Localizes to mitochondria-associated endoplasmic reticulum membrane (MAM); palmitoylation is required for MAM localization (PubMed:22045338, PubMed:27502484, PubMed:31304984).

Tissue Location

Ubiquitous (PubMed:11152479). Highly expressed in kidney, liver, placenta and lung (PubMed:11152479)


TMX1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TMX1 Antibody (C-term) - Images





Anti-TMX1 Antibody (C-term) at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

TMX1 Antibody (C-term) - Background

May participate in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions.