

INTS9 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20367b**Specification**

INTS9 Antibody (C-term) - Product Information

| | |
|-------------------|--|
| Application | WB,E |
| Primary Accession | Q9NV88 |
| Other Accession | Q8K114 , Q4R5Z4 , Q2KJA6 |
| Reactivity | Human |
| Predicted | Bovine, Monkey, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 73814 |
| Antigen Region | 489-515 |

INTS9 Antibody (C-term) - Additional Information**Gene ID** 55756**Other Names**

Integrator complex subunit 9, Int9, Protein related to CPSF subunits of 74 kDa, RC-74, INTS9, RC74

Target/Specificity

This INTS9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 489-515 amino acids from the C-terminal region of human INTS9.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

INTS9 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

INTS9 Antibody (C-term) - Protein Information**Name** INTS9 {ECO:0000303|PubMed:29471365, ECO:0000312|EMBL:BAA91867.1}

Function Component of the integrator complex, a multiprotein complex that terminates RNA polymerase II (Pol II) transcription in the promoter-proximal region of genes (PubMed:[25201415](#), PubMed:[33243860](#), PubMed:[33548203](#), PubMed:[38570683](#)). The integrator complex provides a quality checkpoint during transcription elongation by driving premature transcription termination of transcripts that are unfavorably configured for transcriptional elongation: the complex terminates transcription by (1) catalyzing dephosphorylation of the C-terminal domain (CTD) of Pol II subunit POLR2A/RPB1 and SUPT5H/SPT5, (2) degrading the exiting nascent RNA transcript via endonuclease activity and (3) promoting the release of Pol II from bound DNA (PubMed:[33243860](#), PubMed:[38570683](#)). The integrator complex is also involved in terminating the synthesis of non-coding Pol II transcripts, such as enhancer RNAs (eRNAs), small nuclear RNAs (snRNAs), telomerase RNAs and long non-coding RNAs (lncRNAs) (PubMed:[16239144](#), PubMed:[22252320](#), PubMed:[26308897](#), PubMed:[30737432](#)). Mediates recruitment of cytoplasmic dynein to the nuclear envelope, probably as component of the integrator complex (PubMed:[23904267](#)).


Cellular Location

Nucleus. Cytoplasm


INTS9 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

INTS9 Antibody (C-term) - Images

INTS9 Antibody (C-term) (Cat. #AP20367b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the INTS9 antibody detected the INTS9 protein (arrow).



Anti-INTS9 Antibody (C-term) at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 74 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

INTS9 Antibody (C-term) - Background

Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.