

ZNF12 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20385a**Specification**

ZNF12 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P17014
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	81202
Antigen Region	102-128

ZNF12 Antibody (N-term) - Additional Information**Gene ID** 7559**Other Names**

Zinc finger protein 12, Gonadotropin-inducible ovary transcription repressor 3, GIOT-3, Zinc finger protein 325, Zinc finger protein KOX3, ZNF12, GIOT3, KOX3, ZNF325

Target/Specificity

This ZNF12 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 102-128 amino acids from the N-terminal region of human ZNF12.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZNF12 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF12 Antibody (N-term) - Protein Information**Name** ZNF12**Synonyms** GIOT3, KOX3, ZNF325

Function Transcriptional repressor which suppresses activation protein 1 (AP-1)- and serum response element (SRE)-mediated transcriptional activity.

Cellular Location

Nucleus.

Tissue Location


Widely expressed in various adult tissues and embryonic developmental stages (isoform 3)

ZNF12 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF12 Antibody (N-term) - Images



Anti-ZNF12 Antibody (N-term) at 1:1000 dilution + NCI-H292 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 81 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ZNF12 Antibody (N-term) - Background

Transcriptional repressor which suppresses activation protein 1 (AP-1)-and serum response element (SRE)-mediated transcriptional activity.