

CDCA7L Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20468b

Specification

CDCA7L Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	B3KTR5
Other Accession	Q96GN5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	290-317

CDCA7L Antibody (C-term) - Additional Information

Other Names

CDCA7L;HR1; JPO2; R1; Cell division cycle-associated 7-like protein; Cell division cycle-associated 7-like protein; Protein JPO2; Cell division cycle-associated 7-like protein; Transcription factor RAM2

Target/Specificity

This CDCA7L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 290-317 amino acids from the C-terminal region of human CDCA7L.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDCA7L Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.


CDCA7L Antibody (C-term) - Protein Information

CDCA7L Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDCA7L Antibody (C-term) - Images



CDCA7L Antibody (C-term) (Cat. #AP20468b) western blot analysis in 293,A2058,Ramos,U251,U937 cell line lysates (35ug/lane).This demonstrates the CDCA7L antibody detected the CDCA7L protein (arrow).

CDCA7L Antibody (C-term) - Background

Plays a role in transcriptional regulation as a repressor that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. Plays an important oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. Involved in apoptotic signaling pathways; May act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1.

CDCA7L Antibody (C-term) - References

Huang A., et al. Cancer Res. 65:5607-5619(2005).
Chen K., et al. J. Biol. Chem. 280:11552-11559(2005).
Cathomen T., et al. Submitted (OCT-2002) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).
Scherer S.W., et al. Science 300:767-772(2003).