

R Cask Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20487c

Specification

R Cask Antibody (Center) - Product Information

Application	
Primary Accession	
Other Accession	
Reactivity	
Predicted	
Host	
Clonality	
Isotype	
Calculated MW	
Antigen Region	

WB,E <u>062915</u> <u>070589</u>, <u>014936</u> Rat Human, Mouse Rabbit Polyclonal Rabbit IgG 103259 589-616

R Cask Antibody (Center) - Additional Information

Gene ID 29647

Other Names

Peripheral plasma membrane protein CASK, Calcium/calmodulin-dependent serine protein kinase, Cask

Target/Specificity

This Rat Cask antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 589-616 amino acids from the Central region of rat Cask.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

R Cask Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

R Cask Antibody (Center) - Protein Information

Name Cask {ECO:0000312|RGD:62004}



Function Multidomain scaffolding Mg(2+)-independent protein kinase that catalyzes the phosphotransfer from ATP to proteins such as NRXN1, and plays a role in synaptic transmembrane protein anchoring and ion channel trafficking (By similarity). Multidomain scaffolding protein with a role in synaptic transmembrane protein anchoring and ion channel trafficking.ontributes to neural development and regulation of gene expression via interaction with the transcription factor TBR1. Binds to cell-surface proteins, including amyloid precursor protein, neurexins, and syndecans. May mediate a link between the extracellular matrix and the actin cytoskeleton via its interaction with syndecan and with the actin/spectrin-binding protein 4.1. Component of the LIN-10-LIN-2-LIN-7 complex, which associates with the motor protein KIF17 to transport vesicles containing N-methyl-D-aspartate (NMDA) receptor subunit NR2B along microtubules (By similarity).

Cellular Location

Nucleus. Cytoplasm. Cell membrane; Peripheral membrane protein

Tissue Location

Expressed in the foot process layer of podocytes in the kidney glomerulus and in tubules (at protein level). Detected in brain and neurons.

R Cask Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- R Cask Antibody (Center) Images



Rat Cask Antibody (Center) (Cat. #AP20487c) western blot analysis in rat brain tissue lysates (35ug/lane).This demonstrates the Rat Cask antibody detected the Rat Cask protein (arrow).

R Cask Antibody (Center) - Background



Multidomain scaffolding protein with a role in synaptic transmembrane protein anchoring and ion channel trafficking. Contributes to neural development and regulation of gene expression via interaction with the transcription factor TRB1. Binds to cell-surface proteins, including amyloid precursor protein, neurexins, and syndecans. May mediate a link between the extracellular matrix and the actin cytoskeleton via its interaction with syndecan and with the actin/spectrin-binding protein 4.1.

R Cask Antibody (Center) - References

Hata Y., et al. J. Neurosci. 16:2488-2494(1996). Butz S., et al. Cell 94:773-782(1998). Hsueh Y.P., et al. Nature 404:298-302(2000). Tabuchi K., et al. J. Neurosci. 22:4264-4273(2002). Chetkovich D.M., et al. J. Neurosci. 22:6415-6425(2002).