

**H2AFY2 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20530c****Specification**

---

**H2AFY2 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q9P0M6</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	40058
Antigen Region	148-169

**H2AFY2 Antibody (Center) - Additional Information****Gene ID** 55506**Other Names**

Core histone macro-H2A2, Histone macroH2A2, mH2A2, H2AFY2, MACROH2A2

**Target/Specificity**

This H2AFY2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 148-169 amino acids from the Central region of human H2AFY2.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

H2AFY2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**H2AFY2 Antibody (Center) - Protein Information****Name** MACROH2A2 ([HGNC:14453](#))**Function** Variant histone H2A which replaces conventional H2A in a subset of nucleosomes where it represses transcription. Nucleosomes wrap and compact DNA into chromatin, limiting DNA

accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in stable X chromosome inactivation.

#### Cellular Location


Nucleus. Chromosome. Note=Enriched in inactive X chromosome chromatin (PubMed:11331621, PubMed:11262398) and in senescence- associated heterochromatin (PubMed:15621527)

#### H2AFY2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### H2AFY2 Antibody (Center) - Images



H2AFY2 Antibody (Center) (Cat. #AP20530c) western blot analysis in CEM,NCI-H460 cell line lysates (35ug/lane).This demonstrates the H2AFY2 antibody detected the H2AFY2 protein (arrow).

#### H2AFY2 Antibody (Center) - Background

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes where it represses transcription. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in stable X chromosome inactivation.