

**PIK3R1/2 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20768c****Specification**

---

**PIK3R1/2 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P27986</a>
Other Accession	<a href="#">O00459</a> , <a href="#">P23726</a> , <a href="#">Q63787</a> , <a href="#">P26450</a> , <a href="#">P23727</a> , <a href="#">Q8UUU2</a>
Reactivity	Human, Rat
Predicted	Xenopus, Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	83598

**PIK3R1/2 Antibody (Center) - Additional Information****Gene ID** 5295**Other Names**

Phosphatidylinositol 3-kinase regulatory subunit alpha, PI3-kinase regulatory subunit alpha, PI3K regulatory subunit alpha, PtdIns-3-kinase regulatory subunit alpha, Phosphatidylinositol 3-kinase 85 kDa regulatory subunit alpha, PI3-kinase subunit p85-alpha, PtdIns-3-kinase regulatory subunit p85-alpha, PIK3R1, GRB1

**Target/Specificity**

This PIK3R1/2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 337-370 amino acids from the Central region of human PIK3R1/2.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PIK3R1/2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**PIK3R1/2 Antibody (Center) - Protein Information**

**Name** PIK3R1

**Synonyms** GRB1

**Function** Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed:[17626883](#), PubMed:[19805105](#), PubMed:[7518429](#)). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (PubMed:[20348923](#)).

#### Tissue Location


Isoform 2 is expressed in skeletal muscle and brain, and at lower levels in kidney and cardiac muscle. Isoform 2 and isoform 4 are present in skeletal muscle (at protein level)

#### PIK3R1/2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### PIK3R1/2 Antibody (Center) - Images



Western blot analysis of lysates from U-937, Jurkat, C6 cell line (from left to right), using PIK3R1/2 Antibody (Center)(Cat. #AP20768c). AP20768c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

**PIK3R1/2 Antibody (Center) - Background**

Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling.

**PIK3R1/2 Antibody (Center) - References**

Skolnik E.Y.,et al.Cell 65:83-90(1991).  
Antonetti D.A.,et al.Mol. Cell. Biol. 16:2195-2203(1996).  
Udelhoven M.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Totoki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.