

MOUSE Brd7 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20823c**Specification**

MOUSE Brd7 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O88665
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	387-418

MOUSE Brd7 Antibody (Center) - Additional Information**Gene ID** 26992**Other Names**

Bromodomain-containing protein 7, 75 kDa bromodomain protein, Brd7, Bp75

Target/Specificity

This MOUSE Brd7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 387-418 amino acids from the Central region of human MOUSE Brd7.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MOUSE Brd7 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MOUSE Brd7 Antibody (Center) - Protein Information**Name** Brd7**Synonyms** Bp75**Function** Acts both as coactivator and as corepressor. May play a role in chromatin remodeling.

Transcriptional corepressor that down-regulates the expression of target genes. Binds to target promoters, leading to increased histone H3 acetylation at 'Lys-9' (H3K9ac). Binds to the ESR1 promoter. Recruits BRCA1 and POU2F1 to the ESR1 promoter. Coactivator for TP53-mediated activation of transcription of a set of target genes. Required for TP53-mediated cell-cycle arrest in response to oncogene activation. Promotes acetylation of TP53 at 'Lys-382', and thereby promotes efficient recruitment of TP53 to target promoters. Inhibits cell cycle progression from G1 to S phase (By similarity). Activator of the Wnt signaling pathway in a DVL1-dependent manner by negatively regulating the GSK3B phosphotransferase activity. Induces dephosphorylation of GSK3B at 'Tyr-216'. Down-regulates TRIM24-mediated activation of transcriptional activation by AR.

Cellular Location

Nucleus. Chromosome {ECO:0000250|UniProtKB:Q9NPI1}


Tissue Location

Ubiquitous..


MOUSE Brd7 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MOUSE Brd7 Antibody (Center) - Images

Western blot analysis of lysate from mouse lung tissue lysate, using MOUSE Brd7 Antibody (Center)(Cat. #AP20823c). AP20823c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.



All lanes : Anti-Brd7 Antibody (Center) at 1:2000 dilution Lane 1: mouse heart lysate Lane 2: mouse kidney lysate Lane 3: mouse lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 74 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

MOUSE Brd7 Antibody (Center) - Background

Acts both as coactivator and as corepressor. May play a role in chromatin remodeling. Transcriptional corepressor that down-regulates the expression of target genes. Binds to target promoters, leading to increased histone H3 acetylation at 'Lys-9' (H3K9ac). Binds to the ESR1 promoter. Recruits BRCA1 and POU2F1 to the ESR1 promoter. Coactivator for TP53-mediated activation of transcription of a set of target genes. Required for TP53-mediated cell-cycle arrest in response to oncogene activation. Promotes acetylation of TP53 at 'Lys-382', and thereby promotes efficient recruitment of TP53 to target promoters. Inhibits cell cycle progression from G1 to S phase (By similarity). Activator of the Wnt signaling pathway in a DVL1-dependent manner by negatively regulating the GSK3B phosphotransferase activity. Induces dephosphorylation of GSK3B at 'Tyr-216'. Down-regulates TRIM24- mediated activation of transcriptional activation by AR.

MOUSE Brd7 Antibody (Center) - References

Cuppen E.,et al.FEBS Lett. 459:291-298(1999).
Carninci P.,et al.Science 309:1559-1563(2005).
Kim S.,et al.Cancer Res. 63:4792-4795(2003).
Kaeser M.D.,et al.J. Biol. Chem. 283:32254-32263(2008).
Kikuchi M.,et al.Biochim. Biophys. Acta 1793:1828-1836(2009).