

(Mouse) Sox17 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20861c

Specification

(Mouse) Sox17 Antibody (C-term) - Product Information

Application WB,E
Primary Accession 061473

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 44646

(Mouse) Sox17 Antibody (C-term) - Additional Information

Gene ID 20671

Other Names

Transcription factor SOX-17, Sox17, Sox-17

Target/Specificity

This (Mouse) Sox17 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 376-409 amino acids from the C-terminal region of human (Mouse) Sox17.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(Mouse) Sox17 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

(Mouse) Sox17 Antibody (C-term) - Protein Information

Name Sox17

Synonyms Sox-17



Function Acts as a transcription regulator that binds target promoter DNA and bends the DNA (PubMed:19328208, PubMed:24153254, PubMed:8636240). Binds to the sequences 5'-AACAAT-'3 or 5'-AACAAG-3' (PubMed:8636240). Modulates transcriptional regulation via WNT3A. Inhibits Wnt signaling. Promotes degradation of activated CTNNB1. Plays a key role in the regulation of embryonic development (PubMed:11973269, PubMed:17655922, PubMed:24153254). Required for normal development of the definitive gut endoderm (PubMed:11973269). Required for normal looping of the embryonic heart tube. Plays an important role in embryonic and postnatal vascular development, including development of arteries (PubMed:24153254). Plays an important role in postnatal angiogenesis, where it is functionally redundant with SOX18 (PubMed:16895970). Required for the generation and maintenance of fetal hematopoietic stem cells, and for fetal hematopoiesis (PubMed:17655922). Probable transcriptional activator in the premeiotic germ cells.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267, ECO:0000269|PubMed:20802155, ECO:0000269|PubMed:24153254, ECO:0000269|PubMed:8636240}

Tissue Location


Detected in lung and testis (PubMed:8636240). Detected in endothelial cells around small and large arteries in newborns and adults, but is barely detectable in veins (at protein level) (PubMed:24153254).

(Mouse) Sox17 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

(Mouse) Sox17 Antibody (C-term) - Images



Western blot analysis of lysates from LNCaP, mouse F9 cell line (from left to right), using Sox17



Antibody (C-term)(Cat. #AP20861c). AP20861c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

(Mouse) Sox17 Antibody (C-term) - Background

Acts as transcription regulator that binds target promoter DNA and bends the DNA. Binds to the sequences 5'- AACAAT-'3 or 5'-AACAAAG-3'. Modulates transcriptional regulation via WNT3A. Inhibits Wnt signaling. Promotes degradation of activated CTNNB1. Plays a key role in the regulation of embryonic development. Required for normal looping of the embryonic heart tube. Required for normal development of the definitive gut endoderm. Probable transcriptional activator in the premeiotic germ cells. Isoform 2 (T-SOX17) shows no DNA-binding activity.

(Mouse) Sox17 Antibody (C-term) - References

Kanai Y.,et al.J. Cell Biol. 133:667-681(1996). Carninci P.,et al.Science 309:1559-1563(2005). Layfield R.,et al.Submitted (FEB-1994) to the EMBL/GenBank/DDBJ databases. Kanai-Azuma M.,et al.Development 129:2367-2379(2002). Kim I.,et al.Cell 130:470-483(2007).