

Mouse Frk Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20886c**Specification**

Mouse Frk Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O922K9
Other Accession	Q62662
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	58844

Mouse Frk Antibody (C-term) - Additional Information**Gene ID** 14302**Other Names**

Tyrosine-protein kinase FRK, Beta-cell Src-homology tyrosine kinase, BSK, FYN-related kinase, Intestine tyrosine kinase, Frk, Bsk, Iyk

Target/Specificity

This Mouse Frk antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 482-516 amino acids from the C-terminal region of human Mouse Frk.

Dilution

WB~~1:500-1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Frk Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Frk Antibody (C-term) - Protein Information**Name** Frk**Synonyms** Bsk, Iyk

Function Non-receptor tyrosine-protein kinase that negatively regulates cell proliferation. Positively regulates PTEN protein stability through phosphorylation of PTEN on 'Tyr-336', which in turn prevents its ubiquitination and degradation, possibly by reducing its binding to NEDD4. May function as a tumor suppressor (By similarity).

Cellular Location

Cytoplasm. Nucleus. Note=Predominantly found in the nucleus, with a small fraction found in the cell periphery.

Tissue Location


Expressed in intestinal tract, fetal and adult islets of Langerhans, kidney, liver and lung

Mouse Frk Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Frk Antibody (C-term) - Images



Western blot analysis of lysates from K562 cell line, mouse small intestine, rat stomach tissue lysate(from left to right), using Frk Antibody (C-term)(Cat. #AP20886c). AP20886c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Mouse Frk Antibody (C-term) - Background

Non-receptor tyrosine-protein kinase that negatively regulates cell proliferation. Positively regulates PTEN protein stability through phosphorylation of PTEN on 'Tyr-336', which in turn prevents its ubiquitination and degradation, possibly by reducing its binding to NEDD4. May

function as a tumor suppressor (By similarity).

Mouse Frk Antibody (C-term) - References

Thuveson M.,et al.Biochem. Biophys. Res. Commun. 209:582-589(1995).
Oberg-Welsh C.,et al.Gene 152:239-242(1995).
Carninci P.,et al.Science 309:1559-1563(2005).
Chandrasekharan S.,et al.Mol. Cell. Biol. 22:5235-5247(2002).
Welsh M.,et al.Biochem. J. 382:261-268(2004).