

Mouse Lhx1 Antibody (C-term) Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21094a

Specification

Mouse Lhx1 Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW WB,E <u>P63006</u> <u>P63007</u>, <u>P48742</u> Human Rat Rabbit Polyclonal Rabbit IgG 44780

Mouse Lhx1 Antibody (C-term) - Additional Information

Gene ID 16869

Other Names LIM/homeobox protein Lhx1, LIM homeobox protein 1, Homeobox protein Lim-1, Lhx1, Lim-1, Lim1

Target/Specificity

This Mouse Lhx1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 332-365 amino acids from the C-terminal region of Mouse Lhx1.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Lhx1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Lhx1 Antibody (C-term) - Protein Information

Name Lhx1

Synonyms Lim-1, Lim1



Function Potential transcription factor. May play a role in early mesoderm formation and later in lateral mesoderm differentiation and neurogenesis.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:8793615}

Tissue Location

In mid to late stage embryos, expressed in a restricted region of mesoderm in the primitive streak. At 7.5 days, expressed in a horseshoe shape at the periphery of the node, as well as along both sides of the adjacent notochord. Also present in presumptive lateral and intermediate mesoderm. Later, expression become progressively restricted to intermediate mesoderm, and the developing excretory system including the pronephric region, mesonephros, nephric duct and metanephros. In the metanephros, strongly expressed in renal vesicles and S-shaped and coma-shaped bodies, as well as in the ureteric bud and its derivatives. Also expressed in the dorsal root ganglia. By stage 10.5, also expressed in regions of the central nervous system in the telencephalon through to the spinal cord. In adults, expressed in the cerebellum/medulla and kidney, and at very low levels in the cerebrum.

Mouse Lhx1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Mouse Lhx1 Antibody (C-term) - Images



Western blot analysis of lysate from HepG2 cell line, using Lhx1 Antibody (C-term)(Cat. #AP21094a). AP21094a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Mouse Lhx1 Antibody (C-term) - Background



Potential transcription factor. May play a role in early mesoderm formation and later in lateral mesoderm differentiation and neurogenesis.

Mouse Lhx1 Antibody (C-term) - References

Fujii T.,et al.Dev. Dyn. 199:73-83(1994). Barnes J.D.,et al.Dev. Biol. 161:168-178(1994). Li Y.,et al.Mamm. Genome 10:444-446(1999). Karavanov A.A.,et al.Int. J. Dev. Biol. 40:453-461(1996). Agulnick A.D.,et al.Nature 384:270-272(1996).