

NRBP2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21153a**Specification**

NRBP2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q9NSY0
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	57803

NRBP2 Antibody (C-term) - Additional Information**Gene ID** 340371**Other Names**

Nuclear receptor-binding protein 2, Transformation-related gene 16 protein, TRG-16, NRBP2 (HGNC:19339)

Target/Specificity

This NRBP2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of human NRBP2.

Dilution

WB~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NRBP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NRBP2 Antibody (C-term) - Protein Information**Name** NRBP2 ([HGNC:19339](#))**Function** May regulate apoptosis of neural progenitor cells during their differentiation.


Cellular Location

Cytoplasm.


NRBP2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NRBP2 Antibody (C-term) - Images

Anti-NRBP2 Antibody (C-term) at 1:1000 dilution + mouse liver lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Anti-NRBP2 Antibody (C-term) at 1:500 dilution + A549 whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

NRBP2 Antibody (C-term) - Background

May regulate apoptosis of neural progenitor cells during their differentiation.

NRBP2 Antibody (C-term) - References

Kim J.W.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Wan D.,et al.Proc. Natl. Acad. Sci. U.S.A. 101:15724-15729(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Nusbaum C.,et al.Nature 439:331-335(2006).