

(Mouse) Ephal Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21159a

Specification

(Mouse) Epha1 Antibody (Center) - Product Information

Application WB,E
Primary Accession Q60750
Reactivity Human
Host Rabbit
Clonality polyclonal
Isotype Rabbit IgG
Calculated MW 108578

(Mouse) Epha1 Antibody (Center) - Additional Information

Gene ID 13835

Other Names

Ephrin type-A receptor 1, mEpha1, Embryonic stem cell kinase, Tyrosine-protein kinase receptor ESK, Epha1, Esk

Target/Specificity

This mouse Epha1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 453-487 amino acids from the Central region of mouse Epha1.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(Mouse) Epha1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

(Mouse) Epha1 Antibody (Center) - Protein Information

Name Epha1

Synonyms Esk



Function Receptor tyrosine kinase which binds promiscuously membrane- bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Binds with a low affinity EFNA3 and EFNA4 and with a high affinity to EFNA1 which most probably constitutes its cognate/functional ligand. Upon activation by EFNA1 induces cell attachment to the extracellular matrix inhibiting cell spreading and motility through regulation of ILK and downstream RHOA and RAC. Also plays a role in angiogenesis and regulates cell proliferation. May play a role in apoptosis.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Preferentially expressed in epithelial cells including skin, kidney, liver and thymus (PubMed:11519828, PubMed:18802966). Expressed in myogenic progenitor cells (PubMed:27446912).

(Mouse) Epha1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

(Mouse) Epha1 Antibody (Center) - Images



Anti-(Mouse) Epha1 Antibody (Center) at 1:2000 dilution + A549 whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 109 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

(Mouse) Epha1 Antibody (Center) - Background





Tel: 858.875.1900 Fax: 858.875.1999

Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Binds with a low affinity EFNA3 and EFNA4 and with a high affinity to EFNA1 which most probably constitutes its cognate/functional ligand. Upon activation by EFNA1 induces cell attachment to the extracellular matrix inhibiting cell spreading and motility through regulation of ILK and downstream RHOA and RAC. Plays also a role in angiogenesis and regulates cell proliferation. May play a role in apoptosis.

(Mouse) Epha1 Antibody (Center) - References

Coulthard M.G., et al. Growth Factors 18:303-317(2001). Carninci P., et al. Science 309:1559-1563(2005). Lickliter J.D., et al. Proc. Natl. Acad. Sci. U.S.A. 93:145-150(1996). Duffy S.L., et al. Genesis 46:553-561(2008).