

Mouse Neurog2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21257b

Specification

Mouse Neurog2 Antibody (C-term) - Product Information

Application WB,E
Primary Accession P70447
Reactivity Human
Host Rabbit
Clonality polyclonal
Isotype Rabbit IgG
Calculated MW 28216

Mouse Neurog2 Antibody (C-term) - Additional Information

Gene ID 11924

Other Names

Neurogenin-2, NGN-2, Helix-loop-helix protein mATH-4A, mATH4A, Protein atonal homolog 4, Neurog2, Ath4a, Atoh4, Ngn2

Target/Specificity

This Mouse Neurog2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 247-282 amino acids from the C-terminal region of Mouse Neurog2.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Neurog2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Neurog2 Antibody (C-term) - Protein Information

Name Neurog2

Synonyms Ath4a, Atoh4, Ngn2

Function Transcriptional regulator. Involved in neuronal differentiation. Activates transcription by



binding to the E box (5'- CANNTG-3').

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.

Mouse Neurog2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Neurog2 Antibody (C-term) - Images



Anti-Neurog2 Antibody (C-term)at 1:2000 dilution + Hela whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse Neurog2 Antibody (C-term) - Background

Transcriptional regulator. Involved in neuronal differentiation. Activates transcription by binding to the E box (5'-CANNTG-3').

Mouse Neurog2 Antibody (C-term) - References

Sommer L.,et al.Mol. Cell. Neurosci. 8:221-241(1996). Gradwohl G.,et al.Dev. Biol. 180:227-241(1996). Simmons A.D.,et al.Submitted (SEP-2000) to the EMBL/GenBank/DDBJ databases. Lin C.H.,et al.Dev. Biol. 265:234-245(2004).