

**Mouse Alk Antibody (P1517)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21260a****Specification**

---

**Mouse Alk Antibody (P1517) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P97793</a>
Reactivity	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	174948

**Mouse Alk Antibody (P1517) - Additional Information****Gene ID** 11682**Other Names**

ALK tyrosine kinase receptor, Anaplastic lymphoma kinase, CD246, Alk

**Target/Specificity**

This Mouse Alk antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1517-1550 amino acids Mouse Alk.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Mouse Alk Antibody (P1517) is for research use only and not for use in diagnostic or therapeutic procedures.

**Mouse Alk Antibody (P1517) - Protein Information****Name** Alk {ECO:0000303|PubMed:9053841, ECO:0000312|MGI:MGI:103305}

**Function** Neuronal receptor tyrosine kinase that is essentially and transiently expressed in specific regions of the central and peripheral nervous systems and plays an important role in the genesis and differentiation of the nervous system (PubMed:[15226403](#), PubMed:[16458083](#), PubMed:[16878150](#), PubMed:[19200234](#), PubMed:[30497772](#)). Also acts as a key thinness protein

involved in the resistance to weight gain: in hypothalamic neurons, controls energy expenditure acting as a negative regulator of white adipose tissue lipolysis and sympathetic tone to fine-tune energy homeostasis (PubMed:[32442405](#)). Following activation by ALKAL2 ligand at the cell surface, transduces an extracellular signal into an intracellular response. In contrast, ALKAL1 is not a potent physiological ligand for ALK. Ligand-binding to the extracellular domain induces tyrosine kinase activation, leading to activation of the mitogen-activated protein kinase (MAPK) pathway. Phosphorylates almost exclusively at the first tyrosine of the Y-x-x-x- Y-Y motif. Induces tyrosine phosphorylation of CBL, FRS2, IRS1 and SHC1, as well as of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1. ALK activation may also be regulated by pleiotrophin (PTN) and midkine (MDK). PTN-binding induces MAPK pathway activation, which is important for the anti-apoptotic signaling of PTN and regulation of cell proliferation. MDK-binding induces phosphorylation of the ALK target insulin receptor substrate (IRS1), activates mitogen-activated protein kinases (MAPKs) and PI3-kinase, resulting also in cell proliferation induction. Drives NF-kappa-B activation, probably through IRS1 and the activation of the AKT serine/threonine kinase. Recruitment of IRS1 to activated ALK and the activation of NF-kappa-B are essential for the autocrine growth and survival signaling of MDK (By similarity).

#### Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q9UM73}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9UM73} Note=Membrane attachment is essential for promotion of neuron-like differentiation and cell proliferation arrest through specific activation of the MAP kinase pathway. {ECO:0000250|UniProtKB:Q9UM73}

#### Tissue Location

Mainly expressed in central nervous system (CNS) and other parts of the brain such as the paraventricular nucleus (PVN) of the hypothalamus. Expression is also found in peripheral nervous systems, eye, nasal epithelium, olfactory nerve, tongue, skin, tissue surrounding the esophagus, stomach, midgut, as well as testis and ovary.


### Mouse Alk Antibody (P1517) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Mouse Alk Antibody (P1517) - Images





Anti-Alk Antibody (P1517) at 1:2000 dilution + mouse cerebellum lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 175 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

#### **Mouse Alk Antibody (P1517) - Background**

Neuronal orphan receptor tyrosine kinase that is essentially and transiently expressed in specific regions of the central and peripheral nervous systems and plays an important role in the genesis and differentiation of the nervous system. Transduces signals from ligands at the cell surface, through specific activation of the mitogen-activated protein kinase (MAPK) pathway. Phosphorylates almost exclusively at the first tyrosine of the Y-x-x-x-Y-Y motif. Following activation by ligand, ALK induces tyrosine phosphorylation of CBL, FRS2, IRS1 and SHC1, as well as of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1. Acts as a receptor for ligands pleiotrophin (PTN), a secreted growth factor, and midkine (MDK), a PTN-related factor, thus participating in PTN and MDK signal transduction. PTN-binding induces MAPK pathway activation, which is important for the anti-apoptotic signaling of PTN and regulation of cell proliferation. MDK-binding induces phosphorylation of the ALK target insulin receptor substrate (IRS1), activates mitogen-activated protein kinases (MAPKs) and PI3-kinase, resulting also in cell proliferation induction. Drives NF-kappa-B activation, probably through IRS1 and the activation of the AKT serine/threonine kinase. Recruitment of IRS1 to activated ALK and the activation of NF-kappa-B are essential for the autocrine growth and survival signaling of MDK.

#### **Mouse Alk Antibody (P1517) - References**

- Iwahara T., et al. *Oncogene* 14:439-449(1997).
- Church D.M., et al. *PLoS Biol.* 7:E1000112-E1000112(2009).
- Motegi A., et al. *J. Cell Sci.* 117:3319-3329(2004).
- Vernersson E., et al. *Gene Expr. Patterns* 6:448-461(2006).
- Kuo A.H., et al. *Oncogene* 26:859-869(2007).