

**(Mouse) Jag1 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21264b**

**Specification**

---

**(Mouse) Jag1 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9OXX0</a>
Reactivity	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	134164

**(Mouse) Jag1 Antibody (C-term) - Additional Information**

**Gene ID** 16449

**Other Names**

Protein jagged-1, Jagged1, CD339, Jag1

**Target/Specificity**

This Mouse Jag1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1191-1224 amino acids from the C-terminal region of Mouse Jag1.

**Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

(Mouse) Jag1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**(Mouse) Jag1 Antibody (C-term) - Protein Information**

**Name** Jag1

**Function** Ligand for multiple Notch receptors and involved in the mediation of Notch signaling. May be involved in cell-fate decisions during hematopoiesis. Seems to be involved in early and late stages of mammalian cardiovascular development. Inhibits myoblast differentiation (By similarity).

May regulate fibroblast growth factor-induced angiogenesis.

**Cellular Location**

Membrane; Single-pass type I membrane protein. Cell membrane  
{ECO:0000250|UniProtKB:P78504}


**Tissue Location**

Widely expressed in many tissues, with highest expression in brain, heart, muscle and thymus

**(Mouse) Jag1 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**(Mouse) Jag1 Antibody (C-term) - Images**

Anti-Jag1 Antibody (C-term) at 1:2000 dilution + mouse brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution  
Predicted band size : 134 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

**(Mouse) Jag1 Antibody (C-term) - Background**

Ligand for multiple Notch receptors and involved in the mediation of Notch signaling. May be involved in cell-fate decisions during hematopoiesis. Seems to be involved in early and late stages of mammalian cardiovascular development. Inhibits myoblast differentiation (By similarity). May regulate fibroblast growth factor-induced angiogenesis.

**(Mouse) Jag1 Antibody (C-term) - References**

Shimizu K.,et al.J. Biol. Chem. 274:32961-32969(1999).  
Loomes K.M.,et al.Hum. Mol. Genet. 8:2443-2449(1999).