

(Mouse) Ehmt2 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21266a

Specification

(Mouse) Ehmt2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O9Z148
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	138039

(Mouse) Ehmt2 Antibody (N-term) - Additional Information

Gene ID 110147

Other Names

Histone-lysine N-methyltransferase EHMT2, 211-, Euchromatic histone-lysine N-methyltransferase 2, HLA-B-associated transcript 8, Histone H3-K9 methyltransferase 3, H3-K9-HMTase 3, Protein G9a, Ehmt2, Bat8, G9a, Ng36

Target/Specificity

This Mouse Ehmt2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 227-259 amino acids from the N-terminal region of Mouse Ehmt2.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(Mouse) Ehmt2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

(Mouse) Ehmt2 Antibody (N-term) - Protein Information

Name Ehmt2

Synonyms Bat8, G9a, Ng36

Function Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q96KQ7}. Chromosome {ECO:0000250|UniProtKB:Q96KQ7}. Note=Almost excluded from nucleoli. Associates with euchromatic regions (By similarity). Does not associate with heterochromatin (By similarity) {ECO:0000250|UniProtKB:Q96KQ7}

Tissue Location


Ubiquitous..

(Mouse) Ehmt2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

(Mouse) Ehmt2 Antibody (N-term) - Images



All lanes : Anti-Ehmt2 Antibody (N-term) at 1:2000 dilution Lane 1: NCCIT whole cell lysates Lane

2: HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 138 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

(Mouse) Ehmt2 Antibody (N-term) - Background

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

(Mouse) Ehmt2 Antibody (N-term) - References

Tachibana M., et al. *Genes Dev.* 16:1779-1791(2002).
Xie T., et al. *Genome Res.* 13:2621-2636(2003).
Church D.M., et al. *PLoS Biol.* 7:E1000112-E1000112(2009).
Brown S.E., et al. *Mamm. Genome* 12:916-924(2001).
Tachibana M., et al. *J. Biol. Chem.* 276:25309-25317(2001).