

PPAP2A Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21319b

Specification

PPAP2A Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O14494
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	32156

PPAP2A Antibody (C-term) - Additional Information

Gene ID 8611

Other Names

Lipid phosphate phosphohydrolase 1, PAP2-alpha, Phosphatidate phosphohydrolase type 2a, Phosphatidic acid phosphatase 2a, PAP-2a, PAP2a, PPAP2A, LPP1

Target/Specificity

This PPAP2A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 247-281 amino acids from the C-terminal region of human PPAP2A.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PPAP2A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PPAP2A Antibody (C-term) - Protein Information

Name PLPP1 ([HGNC:9228](#))

Synonyms LPP1, PPAP2A

Function Magnesium-independent phospholipid phosphatase of the plasma membrane that catalyzes the dephosphorylation of a variety of glycerolipid and sphingolipid phosphate esters including phosphatidate/PA, lysophosphatidate/LPA, diacylglycerol pyrophosphate/DGPP, sphingosine 1-phosphate/S1P and ceramide 1- phosphate/C1P (PubMed:[10962286](#), PubMed:[17379599](#), PubMed:[9305923](#), PubMed:[9607309](#), PubMed:[9705349](#)). Also acts on N-oleoyl ethanolamine phosphate/N-(9Z-octadecenoyl)-ethanolamine phosphate, a potential physiological compound (PubMed:[9607309](#)). Through its extracellular phosphatase activity allows both the hydrolysis and the cellular uptake of these bioactive lipid mediators from the milieu, regulating signal transduction in different cellular processes (PubMed:[10962286](#), PubMed:[12909631](#), PubMed:[15461590](#), PubMed:[17379599](#)). It is for instance essential for the extracellular hydrolysis of S1P and subsequent conversion into intracellular S1P (PubMed:[17379599](#)). Involved in the regulation of inflammation, platelets activation, cell proliferation and migration among other processes (PubMed:[12909631](#), PubMed:[15461590](#)). May also have an intracellular activity to regulate phospholipid- mediated signaling pathways (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein Apical cell membrane; Multi-pass membrane protein. Membrane raft; Multi-pass membrane protein. Membrane, caveola {ECO:0000250|UniProtKB:Q61469}; Multi-pass membrane protein

Tissue Location

Widely expressed with highest expression found in prostate (PubMed:9305923). Found to be down-regulated in colon adenocarcinomas (PubMed:9570154). [Isoform 2]: Predominant in heart and pancreas.


PPAP2A Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PPAP2A Antibody (C-term) - Images





Anti-PPAP2A Antibody (C-term) at 1:2000 dilution + human kidney lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-PPAP2A Antibody (C-term) at 1:1000 dilution + human liver lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

PPAP2A Antibody (C-term) - Background

Broad-specificity phosphohydrolase that dephosphorylates exogenous bioactive glycerolipids and sphingolipids. Catalyzes the conversion of phosphatidic acid (PA) to diacylglycerol (DG). Pivotal regulator of lysophosphatidic acid (LPA) signaling in the cardiovascular system. Major enzyme responsible of dephosphorylating LPA in platelets, which terminates signaling actions of LPA. May control circulating, and possibly also regulate localized, LPA levels resulting from platelet activation. It has little activity towards ceramide-1-phosphate (C-1-P) and sphingosine-1-phosphate (S-1-P). The relative catalytic efficiency is LPA > PA > S-1-P > C-1-P. It's down-regulation may contribute to the development of colon adenocarcinoma.

PPAP2A Antibody (C-term) - References

Kai M., et al. J. Biol. Chem. 272:24572-24578(1997).

Leung D.W.,et al.DNA Cell Biol. 17:377-385(1998).
Ulrix W.E.J.,et al.J. Biol. Chem. 273:4660-4665(1998).
Roberts R.,et al.J. Biol. Chem. 273:22059-22067(1998).
Schmutz J.,et al.Nature 431:268-274(2004).