

DPYSL5 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21325b

Specification

DPYSL5 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9BPU6
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	61421

DPYSL5 Antibody (C-term) - Additional Information

Gene ID 56896

Other Names

Dihydropyrimidinase-related protein 5, DRP-5, CRMP3-associated molecule, CRAM, Collapsin response mediator protein 5, CRMP-5, UNC33-like phosphoprotein 6, ULIP-6, DPYSL5, CRMP5, ULIP6

Target/Specificity

This DPYSL5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 521-554 amino acids from the C-terminal region of human DPYSL5.

Dilution

WB~~1:2000

IHC-P~~1:25

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DPYSL5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DPYSL5 Antibody (C-term) - Protein Information

Name DPYSL5

Synonyms CRMP5, ULIP6

Function Involved in the negative regulation of dendrite outgrowth.


Cellular Location
Cytoplasm.

DPYSL5 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DPYSL5 Antibody (C-term) - Images



All lanes : Anti-DPYSL5 Antibody (C-term) at 1:2000 dilution Lane 1: Y79 whole cell lysates Lane 2: SH-SY5Y whole cell lysates Lane 3: Raji whole cell lysates Lane 4: mouse brain lysates Lane 5: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 61 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP21325b staining DPYSL5 in human tonsil sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

DPYSL5 Antibody (C-term) - Background

May have a function in neuronal differentiation and/or axon growth.

DPYSL5 Antibody (C-term) - References

Aguera M., et al. Submitted (MAY-2000) to the EMBL/GenBank/DDBJ databases.
Horiuchi M., et al. FEBS Lett. 480:283-286(2000).
Yu Z., et al. Ann. Neurol. 49:146-154(2001).
Kalnina N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Lubec G., et al. Submitted (DEC-2008) to UniProtKB.