

(DANRE) mtm1 Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21338b

Specification

(DANRE) mtm1 Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	Q32PM3
Reactivity	Zebrafish
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Antigen Region	451-485

(DANRE) mtm1 Antibody (C-Term) - Additional Information

Target/Specificity

This DANRE mtm1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 451-485 amino acids from the region of DANRE mtm1.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(DANRE) mtm1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) mtm1 Antibody (C-Term) - Protein Information


(DANRE) mtm1 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

(DANRE) mtm1 Antibody (C-Term) - Images



Anti-mtm1 Antibody (C-Term) at 1:2000 dilution + ZF4 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 74 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.