

TAB2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21381b**Specification**

TAB2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q9NYJ8
Reactivity	Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	76494

TAB2 Antibody (C-term) - Additional Information**Gene ID** 23118**Other Names**

TGF-beta-activated kinase 1 and MAP3K7-binding protein 2, Mitogen-activated protein kinase kinase kinase 7-interacting protein 2, TAK1-binding protein 2, TAB-2, TGF-beta-activated kinase 1-binding protein 2, TAB2, KIAA0733, MAP3K7IP2

Target/Specificity

This TAB2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 646-680 amino acids from the C-terminal region of human TAB2.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TAB2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TAB2 Antibody (C-term) - Protein Information**Name** TAB2 {ECO:0000303|PubMed:10882101, ECO:0000312|HGNC:HGNC:17075}**Function** Adapter required to activate the JNK and NF-kappa-B signaling pathways through the

specific recognition of 'Lys-63'-linked polyubiquitin chains by its RanBP2-type zinc finger (NZF) (PubMed:[10882101](#), PubMed:[11460167](#), PubMed:[15327770](#), PubMed:[22158122](#), PubMed:[27746020](#), PubMed:[33184450](#), PubMed:[36681779](#)). Acts as an adapter linking MAP3K7/TAK1 and TRAF6 to 'Lys-63'-linked polyubiquitin chains (PubMed:[10882101](#), PubMed:[11460167](#), PubMed:[15327770](#), PubMed:[22158122](#), PubMed:[27746020](#)). The RanBP2-type zinc finger (NZF) specifically recognizes Lys-63'-linked polyubiquitin chains unanchored or anchored to the substrate proteins such as RIPK1/RIP1 and RIPK2: this acts as a scaffold to organize a large signaling complex to promote autophosphorylation of MAP3K7/TAK1, and subsequent activation of I- kappa-B-kinase (IKK) core complex by MAP3K7/TAK1 (PubMed:[15327770](#), PubMed:[18079694](#), PubMed:[22158122](#)). Also recognizes and binds Lys-63'-linked polyubiquitin chains of heterotypic 'Lys-63'-/'Lys-48'-linked branched ubiquitin chains (PubMed:[27746020](#)). Regulates the IL1-mediated translocation of NCOR1 out of the nucleus (By similarity). Involved in heart development (PubMed:[20493459](#)).

Cellular Location

Membrane; Peripheral membrane protein. Endosome membrane; Peripheral membrane protein. Lysosome membrane; Peripheral membrane protein. Cytoplasm, cytosol. Note=Following IL1 stimulation, translocation occurs from the membrane to cytosol (PubMed:10882101) Interaction with TRIM38 promotes translocation from cytosol to endosome and lysosome (PubMed:24434549).

Tissue Location


Widely expressed. In the embryo, expressed in the ventricular trabeculae, endothelial cells of the conotruncal cushions of the outflow tract and in the endothelial cells lining the developing aortic valves.


TAB2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TAB2 Antibody (C-term) - Images





Anti-TAB2 Antibody (C-term) at 1:2000 dilution + C6 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 76 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

TAB2 Antibody (C-term) - Background

Adapter linking MAP3K7/TAK1 and TRAF6. Promotes MAP3K7 activation in the IL1 signaling pathway. The binding of 'Lys-63'- linked polyubiquitin chains to TAB2 promotes autophosphorylation of MAP3K7 at 'Thr-187'. Involved in heart development.

TAB2 Antibody (C-term) - References

Takaesu G., et al. Mol. Cell 5:649-658(2000).
Nagase T., et al. DNA Res. 5:277-286(1998).
Nakajima D., et al. DNA Res. 9:99-106(2002).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Ebert L., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.