

UNC5B Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21410c

Specification

UNC5B Antibody (Center) - Product Information

Application	
Primary Accession	
Reactivity	
Host	
Clonality	
Isotype	
Calculated MW	

WB,E <u>O8IZJ1</u> Mouse, Rat Rabbit polyclonal Rabbit IgG 103638

UNC5B Antibody (Center) - Additional Information

Gene ID 219699

Other Names

Netrin receptor UNC5B, Protein unc-5 homolog 2, Protein unc-5 homolog B, p53-regulated receptor for death and life protein 1, UNC5B, P53RDL1, UNC5H2

Target/Specificity

This UNC5B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 507-541 amino acids from the Central region of human UNC5B.

Dilution WB~~1:2000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UNC5B Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

UNC5B Antibody (Center) - Protein Information

Name UNC5B

Function Receptor for netrin required for axon guidance. Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth



cones may be caused by its association with DCC that may trigger signaling for repulsion (By similarity). Functions as a netrin receptor that negatively regulates vascular branching during angiogenesis. Mediates retraction of tip cell filopodia on endothelial growth cones in response to netrin (By similarity). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (PubMed:<u>12598906</u>). Mediates apoptosis by activating DAPK1. In the absence of NTN1, activates DAPK1 by reducing its autoinhibitory phosphorylation at Ser-308 thereby increasing its catalytic activity (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:O08722} Membrane raft {ECO:0000250|UniProtKB:O08722}. Note=Associated with lipid rafts. {ECO:0000250|UniProtKB:O08722}

Tissue Location

Highly expressed in brain. Also expressed at lower level in developing lung, cartilage, kidney and hematopoietic and immune tissues.

UNC5B Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

UNC5B Antibody (Center) - Images



All lanes : Anti-UNC5B Antibody (Center) at 1:2000 dilution Lane 1: rat cerebellum lysates Lane 2: mouse brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 104 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



UNC5B Antibody (Center) - Background

Receptor for netrin required for axon guidance. Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. Mediates apoptosis by activating DAPK1. In the absence of NTN1, activates DAPK1 by reducing its autoinhibitory phosphorylation at Ser-308 thereby increasing its catalytic activity.

UNC5B Antibody (Center) - References

Komatsuzaki K.,et al.Biochem. Biophys. Res. Commun. 297:898-905(2002). Tanikawa C.,et al.Nat. Cell Biol. 5:216-223(2003). Clark H.F.,et al.Genome Res. 13:2265-2270(2003). Ota T.,et al.Nat. Genet. 36:40-45(2004). Deloukas P.,et al.Nature 429:375-381(2004).