

CDK5R1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21452a**Specification**

CDK5R1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q15078
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	34060

CDK5R1 Antibody (N-term) - Additional Information**Gene ID** 8851**Other Names**

Cyclin-dependent kinase 5 activator 1, CDK5 activator 1, Cyclin-dependent kinase 5 regulatory subunit 1, TPKII regulatory subunit, Cyclin-dependent kinase 5 activator 1, p35, p35, Cyclin-dependent kinase 5 activator 1, p25, p25, Tau protein kinase II 23 kDa subunit, p23, CDK5R1, CDK5R, NCK5A

Target/Specificity

This CDK5R1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 30-64 amino acids from the N-terminal region of human CDK5R1.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDK5R1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDK5R1 Antibody (N-term) - Protein Information**Name** CDK5R1**Synonyms** CDK5R, NCK5A

Function p35 is a neuron specific activator of CDK5. The complex p35/CDK5 is required for neurite outgrowth and cortical lamination. Involved in dendritic spine morphogenesis by mediating the EFNA1-EPHA4 signaling. Activator of TPKII. The complex p35/CDK5 participates in the regulation of the circadian clock by modulating the function of CLOCK protein: phosphorylates CLOCK at 'Thr-451' and 'Thr-461' and regulates the transcriptional activity of the CLOCK-BMAL1 heterodimer in association with altered stability and subcellular distribution.

Cellular Location

[Cyclin-dependent kinase 5 activator 1, p35]: Cell membrane; Lipid-anchor; Cytoplasmic side. Cell projection, neuron projection. Note=In the primary cortical neurons, p35 is present in the peripheries and nerve terminals.

Tissue Location


Brain and neuron specific.

CDK5R1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDK5R1 Antibody (N-term) - Images



All lanes : Anti-CDK5R1 Antibody (N-term) at 1:2000 dilution Lane 1: human cerebellum lysates Lane 2: mouse brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

CDK5R1 Antibody (N-term) - Background

p35 is a neuron specific activator of CDK5. The complex p35/CDK5 is required for neurite outgrowth and cortical lamination. Involved in dendritic spine morphogenesis by mediating the EFNA1-EPHA4 signaling. Activator of TPKII. The complex p35/CDK5 participates in the regulation of the circadian clock by modulating the function of CLOCK protein: phosphorylates CLOCK at 'Thr-451' and 'Thr-461' and regulates the transcriptional activity of the CLOCK-ARNTL/BMAL1 heterodimer in association with altered stability and subcellular distribution.

CDK5R1 Antibody (N-term) - References

Tsai L.-H.,et al.Nature 371:419-423(1994).
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Patrick G.N.,et al.Nature 402:615-622(1999).
Kerokoski P.,et al.Brain Res. Mol. Brain Res. 106:50-56(2002).