

USP19 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2145b

Specification

USP19 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	<u>094966</u>
Other Accession	<u>NP_006668</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1289-1319

USP19 Antibody (C-term) - Additional Information

Gene ID 10869

Other Names

Ubiquitin carboxyl-terminal hydrolase 19, Deubiquitinating enzyme 19, Ubiquitin thioesterase 19, Ubiquitin-specific-processing protease 19, Zinc finger MYND domain-containing protein 9, USP19, KIAA0891, ZMYND9

Target/Specificity

This USP19 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1289-1319 amino acids from the C-terminal region of human USP19.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

USP19 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

USP19 Antibody (C-term) - Protein Information

Name USP19



Synonyms KIAA0891, ZMYND9

Function Deubiguitinating enzyme that regulates the degradation of various proteins by removing ubiquitin moleties, thereby preventing their proteasomal degradation. Stabilizes RNF123, which promotes CDKN1B degradation and contributes to cell proliferation (By similarity). Decreases the levels of ubiquitinated proteins during skeletal muscle formation and acts to repress myogenesis. Modulates transcription of major myofibrillar proteins. Also involved in turnover of endoplasmicreticulum-associated degradation (ERAD) substrates (PubMed: 19465887, PubMed: 24356957). Mechanistically, deubiquitinates and thereby stabilizes several E3 ligases involved in the ERAD pathway including SYVN1 or MARCHF6 (PubMed:24356957). Regulates the stability of other E3 ligases including BIRC2/c-IAP1 and BIRC3/c-IAP2 by preventing their ubiquitination (PubMed:<u>21849505</u>). Required for cells to mount an appropriate response to hypoxia by rescuing HIF1A from degradation in a non-catalytic manner and by mediating the deubiguitination of FUNDC1 (PubMed:22128162, PubMed:33978709). Attenuates mitochondrial damage and ferroptosis by targeting and stabilizing NADPH oxidase 4/NOX4 (PubMed:<u>38943386</u>). Negatively regulates TNF-alpha- and IL-1beta- triggered NF-kappa-B activation by hydrolyzing 'Lys-27'- and 'Lys-63'- linked polyubiguitin chains from MAP3K7 (PubMed:<u>31127032</u>). Modulates also the protein level and aggregation of polyQ-expanded huntingtin/HTT through HSP90AA1 (PubMed: 33094816).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein. Note=Accumulates in the mitochondria-associated ER membrane (MAM) in response to hypoxia

USP19 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

USP19 Antibody (C-term) - Images



Western blot analysis of hUSP19-L1304 (Cat. #AP2145b) in Jurkat cell line lysates (35ug/lane). USP19 (arrow) was detected using the purified Pab.



USP19 Antibody (C-term) - Background

Modification of target proteins by ubiquitin participates in a wide array of biological functions. Proteins destined for degradation or processing via the 26 S proteasome are coupled to multiple copies of ubiquitin. However, attachment of ubiquitin or ubiquitin-related molecules may also result in changes in subcellular distribution or modification of protein activity. An additional level of ubiquitin regulation, deubiquitination, is catalyzed by proteases called deubiquitinating enzymes, which fall into four distinct families. Ubiquitin C-terminal hydrolases, ubiquitin-specific processing proteases (USPs),1 OTU-domain ubiquitin-aldehyde-binding proteins, and Jab1/Pad1/MPN-domain-containing metallo-enzymes. Among these four families, USPs represent the most widespread and represented deubiquitinating enzymes across evolution. USPs tend to release ubiquitin from a conjugated protein. They display similar catalytic domains containing conserved Cys and His boxes but divergent N-terminal and occasionally C-terminal extensions, which are thought to function in substrate recognition, subcellular localization, and protein-protein interactions.

USP19 Antibody (C-term) - References

Nagase, T., et al., DNA Res. 5(6):355-364 (1998).