

AKR7L Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21480c

Specification

AKR7L Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q8NHP1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	36970
Antigen Region	207-237

AKR7L Antibody (Center) - Additional Information

Gene ID 246181

Other Names

Aflatoxin B1 aldehyde reductase member 4, 1---, AFB1 aldehyde reductase 3, AFB1-AR 3, Aldoketoreductase 7-like, AKR7L, AFAR3, AKR7A4

Target/Specificity

This AKR7L antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 207-237 amino acids from the Central region of human AKR7L.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AKR7L Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

AKR7L Antibody (Center) - Protein Information

Name AKR7L

Synonyms AFAR3 {ECO:0000303|PubMed:12879023}, AKR

Function Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen (By similarity).

Tissue Location


Mainly expressed in uterus.

AKR7L Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AKR7L Antibody (Center) - Images



All lanes : Anti-AKR7L Antibody (Center) at 1:2000 dilution Lane 1: HepG2 whole cell lysates Lane 2: HL-60 whole cell lysates Lane 3: HeLa whole cell lysates Lane 4: Jurkat whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-AKR7L Antibody (Center) at 1:1000 dilution + mouse liver lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

AKR7L Antibody (Center) - Background

Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen (By similarity).

AKR7L Antibody (Center) - References

Gregory S.G., et al. Nature 441:315-321(2006).
Prabl C., et al. Oncogene 22:4765-4773(2003).