

RBBP5 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21555b**Specification**

RBBP5 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q15291
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	59153

RBBP5 Antibody (C-term) - Additional Information**Gene ID** 5929**Other Names**

Retinoblastoma-binding protein 5, RBBP-5, Retinoblastoma-binding protein RBQ-3, RBBP5, RBQ3

Target/Specificity

This RBBP5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 505-559 amino acids from the C-terminal region of human RBBP5.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RBBP5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RBBP5 Antibody (C-term) - Protein Information**Name** RBBP5**Synonyms** RBQ3**Function** In embryonic stem (ES) cells, plays a crucial role in the differentiation potential,

particularly along the neural lineage, regulating gene induction and H3 'Lys-4' methylation at key developmental loci, including that mediated by retinoic acid (By similarity). Does not affect ES cell self-renewal (By similarity). Component or associated component of some histone methyltransferase complexes which regulates transcription through recruitment of those complexes to gene promoters (PubMed:[19131338](#)). As part of the MLL1/MLL complex, involved in mono-, di- and trimethylation at 'Lys-4' of histone H3 (PubMed:[19556245](#)). Histone H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation (PubMed:[19556245](#)). In association with ASH2L and WDR5, stimulates the histone methyltransferase activities of KMT2A, KMT2B, KMT2C, KMT2D, SETD1A and SETD1B (PubMed:[21220120](#), PubMed:[22266653](#)).

Cellular Location

Nucleus.


Tissue Location

Ubiquitously expressed.


RBBP5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RBBP5 Antibody (C-term) - Images

Anti-RBBP5 Antibody (C-term) at 1:2000 dilution + K562 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 59 kDa Blocking/Dilution buffer: 5% NFD/MBST.



Anti-RBBP5 Antibody (C-term) at 1:2000 dilution + NIH/3T3 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 59 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

RBBP5 Antibody (C-term) - Background

In embryonic stem (ES) cells, plays a crucial role in the differentiation potential, particularly along the neural lineage, regulating gene induction and H3 'Lys-4' methylation at key developmental loci, including that mediated by retinoic acid (By similarity). As part of the MLL1/MLL complex, involved in mono-, di- and trimethylation at 'Lys-4' of histone H3. Histone H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation.

RBBP5 Antibody (C-term) - References

Saijo M., et al. *Genomics* 27:511-519(1995).
Ota T., et al. *Nat. Genet.* 36:40-45(2004).
Gregory S.G., et al. *Nature* 441:315-321(2006).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Hughes C.M., et al. *Mol. Cell* 13:587-597(2004).