

WIPF3 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21633a**Specification**

WIPF3 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	A6NGB9
Reactivity	Human, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	49458

WIPF3 Antibody (N-term) - Additional Information**Gene ID** 644150**Other Names**

WAS/WASL-interacting protein family member 3, Corticosteroids and regional expression protein 16 homolog, WIPF3, CR16

Target/Specificity

This WIPF3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 109-142 amino acids from the N-terminal region of human WIPF3.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

WIPF3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

WIPF3 Antibody (N-term) - Protein Information**Name** WIPF3**Synonyms** CR16

Function May be a regulator of cytoskeletal organization. May have a role in spermatogenesis (By similarity).

Cellular Location


Cytoplasm. Note=In hippocampal neurons colocalizes with WASL in the cell body, axons and the growth cone

WIPF3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WIPF3 Antibody (N-term) - Images



All lanes : Anti-WIPF3 Antibody (N-term) at 1:2000 dilution Lane 1: human liver lysate Lane 2: rat liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

WIPF3 Antibody (N-term) - Background

May be a regulator of cytoskeletal organization. May have a role in spermatogenesis (By similarity).

WIPF3 Antibody (N-term) - References

Hillier L.W.,et al.Nature 424:157-164(2003).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Ebert L.,et al.Submitted (JAN-2003) to the EMBL/GenBank/DDBJ databases.

Zettl M., et al. Curr. Biol. 12:1617-1622(2002).