

PIK3R2 (Y464) Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21653a

Specification

PIK3R2 (Y464) Antibody - Product Information

Application	WB,E
Primary Accession	<u>000459</u>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG

PIK3R2 (Y464) Antibody - Additional Information

Gene ID 5296

Other Names

Phosphatidylinositol 3-kinase regulatory subunit beta, PI3-kinase regulatory subunit beta, PI3K regulatory subunit beta, PtdIns-3-kinase regulatory subunit beta, Phosphatidylinositol 3-kinase 85 kDa regulatory subunit beta, PI3-kinase subunit p85-beta, PtdIns-3-kinase regulatory subunit p85-beta, PtdIns-3-kinase regulatory subunit

Target/Specificity

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 430-470 amino acids from human PIK3R2.

Dilution WB~~1:2000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PIK3R2 (Y464) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PIK3R2 (Y464) Antibody - Protein Information

Name PIK3R2

Function Regulatory subunit of phosphoinositide-3-kinase (PI3K), a kinase that phosphorylates




PtdIns(4,5)P2 (Phosphatidylinositol 4,5- bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDPK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Binds to activated (phosphorylated) protein-tyrosine kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Indirectly regulates autophagy (PubMed:<u>23604317</u>). Promotes nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin- dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (By similarity).

PIK3R2 (Y464) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PIK3R2 (Y464) Antibody - Images



All lanes : Anti-PIK3R2 Antibody (Y464) at 1:2000 dilution Lane 1: K562 whole cell lysate Lane 2: Raji whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 82 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

PIK3R2 (Y464) Antibody - Background

Regulatory subunit of phosphoinositide-3-kinase (PI3K), a kinase that phosphorylates PtdIns(4,5)P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5- trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDPK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Binds to activated (phosphorylated) protein-tyrosine kinases, through its SH2 domain,



and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Indirectly regulates autophagy (PubMed:23604317). Promotes nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin- dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (By similarity).

PIK3R2 (Y464) Antibody - References

Volinia S., et al.Oncogene 7:789-793(1992). Janssen J.W.G., et al.Oncogene 16:1767-1772(1998). Grimwood J., et al.Nature 428:529-535(2004). Braunger J., et al.Oncogene 14:2619-2631(1997). Igarashi K., et al.Biochem. Biophys. Res. Commun. 246:95-99(1998).