

**VDU1 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2166B****Specification**

---

**VDU1 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8TEY7</a>
Other Accession	<a href="#">A6QNM7</a> , <a href="#">Q8TEY6</a>
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	106727
Antigen Region	799-829

**VDU1 Antibody (C-term) - Additional Information****Gene ID** 23032**Other Names**

Ubiquitin carboxyl-terminal hydrolase 33, Deubiquitinating enzyme 33, Ubiquitin thioesterase 33, Ubiquitin-specific-processing protease 33, VHL-interacting deubiquitinating enzyme 1, hVDU1, USP33, KIAA1097, VDU1

**Target/Specificity**

This VDU1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 799-829 amino acids from the C-terminal region of human VDU1.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

VDU1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**VDU1 Antibody (C-term) - Protein Information**

**Name** USP33

**Synonyms** KIAA1097, VDU1

**Function** Deubiquitinating enzyme involved in various processes such as centrosome duplication, cellular migration and beta-2 adrenergic receptor/ADRB2 recycling. Involved in regulation of centrosome duplication by mediating deubiquitination of CCP110 in S and G2/M phase, leading to stabilize CCP110 during the period which centrioles duplicate and elongate. Involved in cell migration via its interaction with intracellular domain of ROBO1, leading to regulate the Slit signaling. Plays a role in commissural axon guidance cross the ventral midline of the neural tube in a Slit-dependent manner, possibly by mediating the deubiquitination of ROBO1. Acts as a regulator of G- protein coupled receptor (GPCR) signaling by mediating the deubiquitination of beta-arrestins (ARRB1 and ARRB2) and beta-2 adrenergic receptor (ADRB2). Plays a central role in ADRB2 recycling and resensitization after prolonged agonist stimulation by constitutively binding ADRB2, mediating deubiquitination of ADRB2 and inhibiting lysosomal trafficking of ADRB2. Upon dissociation, it is probably transferred to the translocated beta-arrestins, leading to beta-arrestins deubiquitination and disengagement from ADRB2. This suggests the existence of a dynamic exchange between the ADRB2 and beta-arrestins. Deubiquitinates DIO2, thereby regulating thyroid hormone regulation. Mediates deubiquitination of both 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains.

**Cellular Location**

Cytoplasm, perinuclear region. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Associates with centrosomes predominantly in S and G2 phases but less in G1 phase (PubMed:23486064).

**Tissue Location**

Widely expressed..


**VDU1 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**VDU1 Antibody (C-term) - Images**





Western blot analysis of anti-VDU1 Pab (Cat. #AP2166b) in T47D cell line lysate (35ug/lane). VDU1 (arrow) was detected using the purified Pab.

### VDU1 Antibody (C-term) - Background

Type 2 iodothyronine deiodinase (D2) is an integral membrane selenoenzyme that stimulates the pro-hormone thyroxine (T4) and supplies the majority of the 3,5,3'-triiodothyronine (T3) essential for brain development.<sup>1</sup> T4 catalysis accelerates selective conjugation to ubiquitin and thereby renders D2 inactive, a posttranslational feedback mechanism used to maintain acceptable T3 levels.<sup>2,3</sup> Ub-D2 was the first recognized substrate for von Hippel-Lindau protein-interacting (pVHL-interacting) deubiquitinating enzyme-1 (VDU1).<sup>4</sup> VDU proteins colocalize with D2 in the endoplasmic reticulum, and their coexpression provides D2 resistance to degradation. VDU1 expression is substantially upregulated in brown adipocytes by norepinephrine or cold exposure, further amplifying D2 activity. VDU1 and VDU2 are coexpressed with D2 in many human tissues, including brain, heart, and skeletal muscle, suggesting potential roles in neurological development, cardiac function, and energy management, in addition to thermal homeostasis. VDU1- or VDU2-catalyzed deubiquitination recycles inactive Ub-D2 to its active deubiquitinated form, circumventing the proteasomal degradation pathway. Thus, Ub-D2 can be either reactivated or degraded, with the balance between these two processes influenced by VDU activity. VDU1-catalyzed D2 deubiquitination may be an important participant in the adaptive mechanism that regulates thyroid hormone action. The reversible ubiquitination-dependent mechanism regulating D2 activity permits highly responsive control of thyroid hormone activation.<sup>5,6</sup>

### VDU1 Antibody (C-term) - References

- Ota, T., et al., Nat. Genet. 36(1):40-45 (2004).
- Li, Z., et al., J. Biol. Chem. 277(7):4656-4662 (2002).
- Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002).
- Kikuno, R., et al., DNA Res. 6(3):197-205 (1999).