

TFAP2C Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21677c**Specification**

TFAP2C Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q92754
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	49177

TFAP2C Antibody (Center) - Additional Information**Gene ID** 7022**Other Names**

Transcription factor AP-2 gamma, AP2-gamma, Activating enhancer-binding protein 2 gamma, Transcription factor ERF-1, TFAP2C

Target/Specificity

This TFAP2C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 114-147 amino acids from the central region of human TFAP2C.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TFAP2C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

TFAP2C Antibody (Center) - Protein Information**Name** TFAP2C

Function Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important

biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer.

Cellular Location


Nucleus.

TFAP2C Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TFAP2C Antibody (Center) - Images



Anti-TFAP2C Antibody (Center) at 1:2000 dilution + SK-BR-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

TFAP2C Antibody (Center) - Background

Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer.

TFAP2C Antibody (Center) - References

Williamson J.A.,et al.Genomics 35:262-264(1996).
McPherson L.A.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:4342-4347(1997).
Haselton M.D.,et al.Submitted (AUG-2001) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Deloukas P.,et al.Nature 414:865-871(2001).