

Dsk2 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2175a**Specification**

Dsk2 Antibody (N-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	Q9UHD9
Other Accession	Q9OZM0 , Q9JJP9 , Q8R317 , Q9UMX0
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	65696
Antigen Region	20-50

Dsk2 Antibody (N-term) - Additional Information**Gene ID** 29978**Other Names**

Ubiquilin-2, Chap1, DSK2 homolog, Protein linking IAP with cytoskeleton 2, PLIC-2, hPLIC-2, Ubiquitin-like product Chap1/Dsk2, UBQLN2, N4BP4, PLIC2

Target/Specificity

This Dsk2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 20-50 amino acids from the N-terminal region of human Dsk2.

Dilution

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Dsk2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Dsk2 Antibody (N-term) - Protein Information

Name UBQLN2

Synonyms N4BP4, PLIC2

Function Plays an important role in the regulation of different protein degradation mechanisms and pathways including ubiquitin- proteasome system (UPS), autophagy and the endoplasmic reticulum- associated protein degradation (ERAD) pathway. Mediates the proteasomal targeting of misfolded or accumulated proteins for degradation by binding (via UBA domain) to their polyubiquitin chains and by interacting (via ubiquitin-like domain) with the subunits of the proteasome (PubMed:[10983987](#)). Plays a role in the ERAD pathway via its interaction with ER-localized proteins FAF2/UBXD8 and HERPUD1 and may form a link between the polyubiquitinated ERAD substrates and the proteasome (PubMed:[18307982](#), PubMed:[24215460](#)). Involved in the regulation of macroautophagy and autophagosome formation; required for maturation of autophagy-related protein LC3 from the cytosolic form LC3-I to the membrane-bound form LC3-II and may assist in the maturation of autophagosomes to autolysosomes by mediating autophagosome-lysosome fusion (PubMed:[19148225](#), PubMed:[20529957](#)). Negatively regulates the endocytosis of GPCR receptors: AVPR2 and ADRB2, by specifically reducing the rate at which receptor-arrestin complexes concentrate in clathrin-coated pits (CCPs) (PubMed:[18199683](#)).

Cellular Location


Cytoplasm. Nucleus. Membrane {ECO:0000250|UniProtKB:Q9QZM0} Cytoplasmic vesicle, autophagosome Note=Colocalizes with a subset of proteasomes, namely those that are cytoskeleton associated or free in the cytosol. Associated with fibers in mitotic cells.


Dsk2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dsk2 Antibody (N-term) - Images






The anti-Dsk2 Pab (Cat. #AP2175a) is used in Western blot to detect Dsk2 in HeLa cell lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with Dsk2 Antibody (N-term) (Cat.#AP2175a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Dsk2 Antibody (N-term) - Background

Dsk2 increases the half-life of proteins destined to be degraded by the proteasome, and may modulate proteasome-mediated protein degradation. The Dsk2 protein binds UBE3A and BTRC, and interacts with the 19S proteasome subunit. In the cytoplasm, Dsk2 colocalizes with the proteasome; it is also associated with fibers in mitotic cells in the nucleus. Dsk2 is highly expressed in mitotic cells from metaphase to telophase, while expression in non-mitotic cells is very low.

Dsk2 Antibody (N-term) - References

Walters, K.J., et al., Biochemistry 41(6):1767-1777 (2002).
Kleijnen, M.F., et al., Mol. Cell 6(2):409-419 (2000).
Ueki, N., et al., Nat. Biotechnol. 16(13):1338-1342 (1998).