

TRIM55 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21795c**Specification**

TRIM55 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9BYV6
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	60466

TRIM55 Antibody (Center) - Additional Information**Gene ID** 84675**Other Names**

Tripartite motif-containing protein 55, Muscle-specific RING finger protein 2, MuRF-2, MuRF2, RING finger protein 29, TRIM55, MURF2, RNF29

Target/Specificity

This TRIM55 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 295-329 amino acids from the Central region of human TRIM55.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TRIM55 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

TRIM55 Antibody (Center) - Protein Information**Name** TRIM55**Synonyms** MURF2, RNF29

Function E3 ubiquitin ligase that plays an important role in regulating cardiac development and contractility, muscle growth, metabolism, and fiber-type differentiation. Acts as a critical factor that regulates cardiomyocyte size during development in concert with TRIM63 by regulating E2F1-mediated gene expression (By similarity). Plays a role in apoptosis induction in cardiomyocytes by promoting ubiquitination of the DUSP1 phosphatase. Promotes non-canonical NF- κ B signaling and B-cell-mediated immune responses by mediating NFKB2 'Lys-48'-linked ubiquitination and processing. In turn, NFKB2 is further processed by valosin-containing protein/VCP, an ATPase that mediates ubiquitin-dependent protein degradation by the proteasome. May play a role in preventing macrophages from producing inflammatory factors and migrating by downregulating the level of nuclear NF-kappa-B subunit RELA. Also modifies PPARG via polyubiquitination and accelerates PPARG proteasomal degradation to inhibit its activity (PubMed:[36737649](#)).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:G3X8Y1}. Cytoplasm {ECO:0000250|UniProtKB:G3X8Y1}. Note=TLR4 signaling pathway promotes nuclear translocation. {ECO:0000250|UniProtKB:G3X8Y1}

Tissue Location


Highly expressed in muscle. Low-level expression in liver.

TRIM55 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRIM55 Antibody (Center) - Images



Anti-TRIM55 Antibody (Center) at 1:2000 dilution + human liver lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.

Predicted band size : 60 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

TRIM55 Antibody (Center) - Background

May regulate gene expression and protein turnover in muscle cells.

TRIM55 Antibody (Center) - References

Centner T.,et al.J. Mol. Biol. 306:717-726(2001).

Pizon V.,et al.J. Cell Sci. 115:4469-4482(2002).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.