

ZBTB7C Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21799c**Specification**

ZBTB7C Antibody (Center) - Product Information

Application	WB,E
Primary Accession	A1YPRO
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	69017

ZBTB7C Antibody (Center) - Additional Information**Gene ID** 201501**Other Names**

Zinc finger and BTB domain-containing protein 7C, Affected by papillomavirus DNA integration in ME180 cells protein 1, APM-1, Zinc finger and BTB domain-containing protein 36, Zinc finger protein 857C, ZBTB7C, APM1, ZBTB36, ZNF857C

Target/Specificity

This ZBTB7C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 264-294 amino acids from the Central region of human ZBTB7C.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZBTB7C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ZBTB7C Antibody (Center) - Protein Information**Name** ZBTB7C**Synonyms** APM1, ZBTB36, ZNF857C

Function May be a tumor suppressor gene.

Tissue Location


Detected in normal cervical keratinocytes, and in some cervical carcinoma cell lines.

ZBTB7C Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZBTB7C Antibody (Center) - Images



Anti-ZBTB7C Antibody (Center) at 1:2000 dilution + Raji whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 69 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ZBTB7C Antibody (Center) - Background

May be a tumor suppressor gene.

ZBTB7C Antibody (Center) - References

Reuter S., et al. EMBO J. 17:215-222(1998).

Gilling M., et al. Submitted (NOV-2006) to the EMBL/GenBank/DDBJ databases.