

**DVL1L1 Antibody (N-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21836a**

### Specification

#### DVL1L1 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	<a href="#">P54792</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	73254
Antigen Region	39-71

#### DVL1L1 Antibody (N-Term) - Additional Information

##### Other Names

Putative segment polarity protein dishevelled homolog DVL1P1, DSH homolog 1-like, Segment polarity protein dishevelled homolog DVL-1-like, Dishevelled-1-like, DVL1P1, DVL, DVL1, DVL1L1

##### Target/Specificity

This DVL1L1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 39-71 amino acids from human DVL1L1.

##### Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

DVL1L1 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### DVL1L1 Antibody (N-Term) - Protein Information

**Name** DVL1P1

**Synonyms** DVL, DVL1, DVL1L1

**Function** May play a role in the signal transduction pathway mediated by multiple Wnt genes.

**Cellular Location**

Cytoplasm.


**Tissue Location**

Expressed in thymus, heart, liver, kidney, brain, skeletal muscle, and pancreas.

**DVL1L1 Antibody (N-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DVL1L1 Antibody (N-Term) - Images**

Anti-DVL1L1 Antibody (N-Term) at 1:2000 dilution + human heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 73 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

**DVL1L1 Antibody (N-Term) - Background**

May play a role in the signal transduction pathway mediated by multiple Wnt genes.

**DVL1L1 Antibody (N-Term) - References**

Pizzuti A.,et al.Am. J. Hum. Genet. 58:722-729(1996).