

GPSM2 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21850a**Specification**

GPSM2 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	P81274
Other Accession	Q8VDU0
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	76662

GPSM2 Antibody (N-Term) - Additional Information**Gene ID** 29899**Other Names**

G-protein-signaling modulator 2, Mosaic protein LGN, GPSM2, LGN

Target/Specificity

This GPSM2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 113-143 amino acids from human GPSM2.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GPSM2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

GPSM2 Antibody (N-Term) - Protein Information**Name** GPSM2**Synonyms** LGN

Function Plays an important role in mitotic spindle pole organization via its interaction with NUMA1 (PubMed:[11781568](#), PubMed:[15632202](#), PubMed:[21816348](#)). Required for cortical dynein-dynactin complex recruitment during metaphase (PubMed:[22327364](#)). Plays a role in metaphase spindle orientation (PubMed:[22327364](#)). Also plays an important role in asymmetric cell divisions (PubMed:[21816348](#)). Has guanine nucleotide dissociation inhibitor (GDI) activity towards G(i) alpha proteins, such as GNAI1 and GNAI3, and thereby regulates their activity (By similarity).

Cellular Location

Cytoplasm. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton, spindle pole. Lateral cell membrane. Note=Localizes in the cytoplasm during interphase and at cell cortex during metaphase (PubMed:[11781568](#), PubMed:[15632202](#), PubMed:[22074847](#)). Colocalizes with NUMA1 to mitotic spindle poles (PubMed:[11781568](#), PubMed:[21816348](#)). Localized at the central and lateral cell cortex regions in a RanGTP-dependent manner (PubMed:[22327364](#)). In horizontally retinal progenitor dividing cells, localized to the lateral cortical region. In vertically retinal progenitor dividing cells, localized at the polar cortical region (By similarity).
{ECO:0000250|UniProtKB:Q8VDU0, ECO:0000269|PubMed:[11781568](#), ECO:0000269|PubMed:[15632202](#), ECO:0000269|PubMed:[21816348](#), ECO:0000269|PubMed:[22074847](#), ECO:0000269|PubMed:[22327364](#)}

Tissue Location


Ubiquitously expressed.

GPSM2 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GPSM2 Antibody (N-Term) - Images



All lanes : Anti-GPSM2 Antibody (N-Term) at 1:2000 dilution Lane 1: U-2OS whole cell lysate Lane 2: A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 77 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

GPSM2 Antibody (N-Term) - Background

Plays an important role in spindle pole orientation. Interacts and contributes to the functional activity of G(i) alpha proteins. Acts to stabilize the apical complex during neuroblast divisions.

GPSM2 Antibody (N-Term) - References

Mochizuki N.,et al.Gene 181:39-43(1996).
Katagiri T.,et al.Submitted (JUL-2008) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Puhl H.L. III,et al.Submitted (JUL-2002) to the EMBL/GenBank/DDBJ databases.