

**NPRL3 Antibody (N-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21877a****Specification**

---

**NPRL3 Antibody (N-Term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q12980</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	63605

**NPRL3 Antibody (N-Term) - Additional Information****Gene ID** 8131**Other Names**

Nitrogen permease regulator 3-like protein, -14 gene protein, Alpha-globin regulatory element-containing gene protein, Protein CGTHBA, NPRL3, C16orf35, CGTHBA, MARE

**Target/Specificity**

This NPRL3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 86-117 amino acids from human NPRL3.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NPRL3 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**NPRL3 Antibody (N-Term) - Protein Information****Name** NPRL3 {ECO:0000303|PubMed:26505888, ECO:0000312|HGNC:HGNC:14124}**Function** As a component of the GATOR1 complex functions as an inhibitor of the amino acid-sensing branch of the mTORC1 pathway (PubMed:[23723238](#), PubMed:[29590090](#), PubMed:[35338845](#)). In response to amino acid depletion, the GATOR1 complex has GTPase

activating protein (GAP) activity and strongly increases GTP hydrolysis by RagA/RRAGA (or RagB/RRAGB) within heterodimeric Rag complexes, thereby turning them into their inactive GDP-bound form, releasing mTORC1 from lysosomal surface and inhibiting mTORC1 signaling (PubMed:[23723238](#), PubMed:[29590090](#), PubMed:[35338845](#)). In the presence of abundant amino acids, the GATOR1 complex is negatively regulated by GATOR2, the other GATOR subcomplex, in this amino acid-sensing branch of the TORC1 pathway (PubMed:[23723238](#)).

#### Cellular Location

Lysosome membrane. Note=Localization to lysosomes is mediated by the KICSTOR complex and is amino acid-independent.

#### Tissue Location


Widely expressed. Expressed in the frontal lobe cortex as well as in the temporal, parietal, and occipital lobes (PubMed:[27173016](#), PubMed:[26505888](#)).

### NPRL3 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NPRL3 Antibody (N-Term) - Images



Anti-NPRL3 Antibody (N-Term) at 1:2000 dilution + HT-1080 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### NPRL3 Antibody (N-Term) - Background

As a component of the GATOR1 complex, inhibitor of the amino acid-sensing branch of the TORC1

pathway. The GATOR1 complex strongly increases GTP hydrolysis by RRAGA and RRAGB within RRAGC- containing heterodimers, thereby deactivating RRAGs, releasing mTORC1 from lysosomal surface and inhibiting mTORC1 signaling.

#### **NPRL3 Antibody (N-Term) - References**

Vyas P.,et al.Genomics 29:679-689(1995).

De Gobbi M.,et al.Science 312:1215-1217(2006).

Martin J.,et al.Nature 432:988-994(2004).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Neklesa T.K.,et al.PLoS Genet. 5:E1000515-E1000515(2009).